

AMI Report: Stanford University Hospital - Model A

year	Statewide Death Rate (%)	Number of Cases Included	Number of Observed Deaths	Number of Expected Deaths	Observed Death Rate (%)	Expected Death Rate (%)	Risk-Adjusted Death Rate (%)	Risk Adjusted Death Rate 95% Confidence Lower Bound	Risk Adjusted Death Rate 95% Confidence Upper Bound	Probability This Rate Occurred by Chance
1991	15.1	182	23	27.9	12.6	15.3	12.4	7.6	17.3	0.171
1992	14.5	173	15	22.5	8.7	13.0	9.6	4.4	14.8	0.040
1993	13.9	178	14	23.8	7.9	13.3	8.2	3.3	13.1	0.011
1994	13.0	191	15	25.2	7.9	13.2	7.7	3.3	12.1	0.010
1995	12.8	198	26	28.0	13.1	14.1	11.9	7.8	16.0	0.379
1996	12.2	177	24	22.8	13.6	12.9	12.8	8.4	17.3	0.421
1997	12.0	183	16	22.4	8.7	12.3	8.6	4.2	12.9	0.073
1998	12.1	215	26	27.9	12.1	13.0	11.2	7.3	15.2	0.389

Model A is a conservative model with risk factors of chronic and personal characters

AMI Report: Stanford University Hospital - Model B

year	Statewide Death Rate (%)	Number of Cases Included	Number of Observed Deaths	Number of Expected Deaths	Observed Death Rate (%)	Expected Death Rate (%)	Risk-Adjusted Death Rate (%)	Risk Adjusted Death Rate 95% Confidence Lower Bound	Risk Adjusted Death Rate 95% Confidence Upper Bound	Probability This Rate Occurred by Chance
1991	15.1	180	23	31.2	12.8	17.3	11.1	7.0	15.2	0.034
1992	14.5	172	15	23.5	8.7	13.6	9.3	4.4	14.1	0.019
1993	13.9	174	14	24.6	8.0	14.1	7.9	3.5	12.3	0.004
1994	12.9	191	15	23.6	7.9	12.3	8.2	3.8	12.6	0.020
1995	12.8	197	26	28.7	13.2	14.5	11.6	7.9	15.3	0.310
1996	12.2	175	24	22.8	13.7	13.0	12.8	8.8	16.9	0.420
1997	12.0	181	16	22.7	8.8	12.5	8.5	4.5	12.4	0.048
1998	12.1	214	26	29.7	12.1	13.9	10.6	7.1	14.1	0.236

Model B is a comprehensive model with more acute risk factors added in