

## AMI Report: Stanislaus Medical Center - Model A

year	Statewide Death Rate (%)	Number of Cases Included	Number of Observed Deaths	Number of Expected Deaths	Observed Death Rate (%)	Expected Death Rate (%)	Risk-Adjusted Death Rate (%)	Risk Adjusted Death Rate 95% Confidence Lower Bound	Risk Adjusted Death Rate 95% Confidence Upper Bound	Probability This Rate Occurred by Chance
1991	15.1	50	8	7.0	16.0	14.1	17.2	7.4	26.9	0.404
1992	14.5	52	6	5.3	11.5	10.2	16.4	5.4	27.4	0.439
1993	13.9	57	5	6.3	8.8	11.0	11.1	1.4	20.7	0.376
1994	13.0	61	6	6.3	9.8	10.3	12.4	3.5	21.4	0.563
1995	12.8	64	8	7.2	12.5	11.3	14.2	5.9	22.4	0.439
1996	12.2	52	7	5.5	13.5	10.6	15.5	6.3	24.6	0.305
1997	12.0	42	4	4.9	9.5	11.6	9.8	0.5	19.2	0.441

*Model A is a conservative model with risk factors of chronic and personal characters*

## AMI Report: Stanislaus Medical Center - Model B

year	Statewide Death Rate (%)	Number of Cases Included	Number of Observed Deaths	Number of Expected Deaths	Observed Death Rate (%)	Expected Death Rate (%)	Risk-Adjusted Death Rate (%)	Risk Adjusted Death Rate 95% Confidence Lower Bound	Risk Adjusted Death Rate 95% Confidence Upper Bound	Probability This Rate Occurred by Chance
1991	15.1	50	8	8.1	16.0	16.1	14.9	6.8	23.1	0.590
1992	14.5	52	6	4.8	11.5	9.2	18.2	7.4	29.0	0.328
1993	13.9	57	5	6.0	8.8	10.6	11.5	2.4	20.7	0.412
1994	12.9	60	6	6.7	10.0	11.1	11.7	3.9	19.4	0.485
1995	12.8	64	8	6.9	12.5	10.8	14.9	7.1	22.7	0.373
1996	12.2	52	7	5.2	13.5	10.0	16.4	7.5	25.4	0.241
1997	12.0	42	4	5.2	9.5	12.4	9.3	1.0	17.5	0.363

*Model B is a comprehensive model with more acute risk factors added in*