

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
Alameda	Alameda County Medical Center – Highland Campus				13.6% (3; 40)		0% (0; 7)	18.5% (5; 30)	4.5% (6; 314)	<b>14.8%</b> (26; 179) Worse	2.4% (3; 152)	6.5% (1; 13)	4.2% (5; 219)
	Alameda Hospital							4.2% (4; 49)	2.9% (7; 194)	12.6% (6; 58)	3.7% (3; 101)	1.9% (1; 37)	2.0% (3; 140)
	Alta Bates Summit Medical Center – Alta Bates Campus				6.6% (7; 68)	1.7% (3; 100)	<b>12.9%</b> (2; 25) Worse	5.4% (7; 107)	2.2% (9; 329)	8.2% (31; 284)	2.7% (7; 217)	1.5% (2; 99)	3.3% (12; 288)
	Alta Bates Summit Medical Center – Summit Campus – Hawthorne	8.2% (1; 5)	0% (0; 7)	12.5% (1; 14)		2.0% (15; 1,079)	0% (0; 18)	<b>4.6%</b> (20; 719) Better	2.0% (13; 487)	7.8% (22; 303)	<b>0.3%</b> (1; 285) Better	1.9% (4; 108)	<b>2.5%</b> (12; 359) Better
	Eden Medical Center				6.7% (11; 75)		0% (0; 8)	4.0% (5; 46)	2.8% (11; 231)	9.5% (42; 227)	<b>0%</b> (0; 129) Better	1.9% (2; 68)	2.5% (8; 212)
	Kaiser Foundation Hospital – Hayward		0% (0; 7)	0% (0; 29)			0% (0; 36)	4.8% (13; 175)	3.1% (20; 503)	<b>7.3%</b> (23; 277) Better	1.0% (4; 343)	2.4% (5; 137)	4.0% (27; 518)
	Kaiser Foundation Hospital – Oakland Campus		12.0% (1; 9)	3.9% (1; 25)			0% (0; 19)	6.1% (14; 152)	2.7% (17; 496)	<b>7.1%</b> (18; 309) Better	2.3% (5; 219)	2.3% (4; 144)	6.3% (34; 414)
	Saint Rose Hospital					<b>3.7%</b> (9; 109) Worse	0% (0; 3)	11.5% (16; 163)	4.2% (7; 220)	7.3% (5; 92)	1.9% (2; 132)	0% (0; 34)	3.7% (3; 135)
	San Leandro Hospital			0% (0; 9)			0% (0; 28)	4.3% (3; 33)	2.1% (8; 339)	<b>3.7%</b> (3; 85) Better	1.6% (2; 119)	3.5% (2; 44)	3.6% (6; 144)
	Valleycare Medical Center		0% (0; 3)	0% (0; 5)		3.4% (5; 180)	0% (0; 21)	6.2% (11; 180)	4.0% (10; 187)	7.4% (3; 71)	3.0% (4; 120)	1.1% (1; 58)	3.9% (11; 264)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

### Definition of Terms:

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurysm Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Washington Hospital – Fremont			0% (0; 4)	3.7% (3; 42)	2.6% (12; 419)	3.9% (1; 20)	6.3% (18; 263)	2.0% (11; 417)	<b>7.5%</b> (30; 308) Better	1.2% (3; 257)	0.8% (1; 99)	4.7% (22; 365)
Amador	Sutter Amador Hospital							<b>22.1%</b> (5; 9) Worse	1.0% (1; 83)	8.9% (4; 34)	<b>18.4%</b> (8; 52) Worse	1.6% (1; 47)	4.7% (6; 127)
Butte	Biggs Gridley Memorial Hospital								11.5% (2; 28)	<b>43.4%</b> (5; 20) Worse	13.7% (1; 12)	0% (0; 11)	8.3% (4; 77)
	Enloe Medical Center – Esplanade Campus			3.2% (1; 28)	2.3% (1; 49)	<b>3.7%</b> (10; 364) Worse	0% (0; 79)	10.7% (22; 266)	3.6% (15; 324)	9.9% (24; 228)	2.8% (8; 242)	2.8% (5; 148)	6.7% (23; 460)
	Feather River Hospital						10.5% (1; 21)	7.7% (8; 49)	<b>0.9%</b> (2; 111) Better	9.2% (8; 83)	1.2% (2; 100)	1.0% (1; 75)	<b>1.3%</b> (3; 137) Better
	Oroville Hospital						0% (0; 16)	4.3% (4; 46)	1.6% (5; 210)	<b>3.6%</b> (5; 120) Better	<b>0%</b> (0; 145) Better	5.4% (3; 54)	<b>1.7%</b> (9; 363) Better
Calaveras	Mark Twain Saint Joseph’s Hospital							19.0% (2; 7)	6.0% (4; 78)	<b>23.9%</b> (6; 32) Worse	2.2% (1; 39)	0% (0; 20)	5.4% (2; 67)
Colusa	Colusa Regional Medical Center							31.9% (1; 3)	4.5% (3; 44)	11.2% (1; 16)	0% (0; 17)		4.9% (3; 58)
Contra Costa	Contra Costa Regional Medical Center				0% (0; 3)			<b>30.0%</b> (4; 19) Worse	2.1% (4; 237)	12.1% (4; 59)	1.2% (1; 76)	0% (0; 20)	2.4% (4; 217)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurysm Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Doctors Medical Center – San Pablo					1.7% (2; 190)	0% (0; 4)	5.9% (11; 178)	3.7% (16; 421)	9.7% (10; 139)	2.8% (5; 163)	1.1% (1; 69)	5.9% (15; 235)
	John Muir Medical Center – Concord Campus	0% (0; 5)		0% (0; 37)		2.2% (8; 274)	0% (0; 48)	8.1% (27; 329)	2.3% (13; 328)	9.6% (17; 152)	3.7% (12; 195)	2.1% (2; 63)	3.7% (12; 257)
	John Muir Medical Center – Walnut Creek Campus			0% (0; 3)	4.3% (11; 110)	1.0% (2; 191)	0% (0; 50)	<b>3.6%</b> (10; 159) <b>Better</b>	2.0% (10; 313)	9.2% (82; 403)	2.2% (9; 261)	1.0% (3; 175)	<b>2.4%</b> (11; 277) <b>Better</b>
	Kaiser Foundation Hospital – Antioch		0% (0; 5)					4.1% (5; 82)	2.8% (10; 258)	9.8% (16; 137)	3.3% (5; 138)	1.5% (1; 72)	4.0% (15; 291)
	Kaiser Foundation Hospital – Walnut Creek ( <a href="#">View Comment Letter</a> )	12.9% (2; 18)	0% (0; 14)	0% (0; 36)		<b>4.2%</b> (6; 54) <b>Worse</b>	0% (0; 49)	6.6% (11; 132)	3.3% (17; 382)	7.6% (14; 218)	0.5% (1; 247)	1.7% (4; 220)	5.4% (27; 412)
	San Ramon Regional Medical Center	0% (0; 3)				2.8% (2; 54)	0% (0; 7)	6.7% (4; 60)	4.5% (5; 94)	4.1% (2; 51)	2.0% (1; 74)	1.6% (1; 30)	2.8% (5; 142)
	Sutter Delta Medical Center					5.2% (3; 53)	0% (0; 16)	6.8% (9; 112)	<b>1.3%</b> (5; 254) <b>Better</b>	7.4% (5; 107)	2.4% (5; 176)	2.8% (3; 71)	4.1% (13; 244)
Del Norte	Sutter Coast Hospital							13.4% (5; 15)	5.0% (6; 79)	11.6% (9; 57)	0.9% (1; 81)	0% (0; 4)	<b>1.8%</b> (3; 92) <b>Better</b>
El Dorado	Barton Memorial Hospital						0% (0; 5)	2.5% (1; 12)	3.7% (2; 66)	13.2% (2; 15)	4.2% (1; 32)	5.3% (1; 28)	0% (0; 73)
	Marshall Medical Center						0% (0; 6)	10.6% (6; 25)	2.3% (7; 172)	<b>5.8%</b> (7; 94) <b>Better</b>	3.9% (4; 94)	1.0% (1; 63)	3.2% (9; 204)
Fresno	Clovis Community Medical Center**					5.3% (1; 36)		4.2% (4; 87)	2.7% (4; 135)	10.2% (4; 73)	1.1% (1; 120)	5.1% (3; 77)	5.5% (12; 288)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	5.8% (29; 500)	5.2% (38; 732)	1.9% (50; 2,567)	6.7% (938; 13,926)	1.9% (933; 48,689)	0.7% (49; 7,411)	7.1% (3,151; 44,510)	3.3% (2,832; 85,934)	10.4% (5,285; 50,988)	2.3% (1,114; 47,720)	2.4% (571; 23,514)	4.6% (3,703; 79,958)
	Coalinga Regional Medical Center								3.3% (1; 37)	44.2% (2; 9)	0% (0; 8)		3.0% (1; 152)
	Community Regional Medical Center – Fresno		29.1% (1; 3)	0% (0; 13)	<b>11.6%</b> (25; 204) Worse	2.0% (9; 430)	0% (0; 14)	7.4% (41; 613)	2.8% (19; 715)	9.8% (97; 643)	2.3% (13; 499)	4.5% (7; 147)	4.8% (32; 756)
	Fresno Heart and Surgical Hospital	62.5% (1; 5)	0% (0; 7)	0% (0; 17)		0% (0; 571)	1.0% (1; 180)	5.9% (4; 117)	4.0% (5; 107)	0% (0; 4)	7.4% (1; 19)		0% (0; 8)
	Kaiser Foundation Hospital – Fresno			0% (0; 15)		0% (0; 3)	3.0% (1; 52)	<b>3.2%</b> (5; 146) Better	2.0% (8; 331)	6.4% (10; 144)	3.3% (4; 115)	4.2% (6; 120)	5.7% (22; 301)
	Saint Agnes Medical Center	0% (0; 3)	0% (0; 5)	0% (0; 7)	7.1% (4; 93)	<b>3.7%</b> (20; 671) Worse	0% (0; 51)	8.2% (39; 518)	<b>4.5%</b> (54; 923) Worse	10.6% (44; 444)	1.4% (8; 511)	2.2% (6; 268)	<b>9.1%</b> (77; 833) Worse
	San Joaquin Valley Rehabilitation Hospital											0% (0; 3)	
	Sierra Kings District Hospital								5.6% (2; 40)	17.1% (1; 12)	0% (0; 6)		6.0% (4; 106)
Glenn	Glenn Medical Center								0% (0; 10)		0% (0; 7)		14.6% (4; 35)
Humboldt	Jerold Phelps Community Hospital								0% (0; 4)				0% (0; 8)
	Mad River Community Hospital						0% (0; 3)	4.6% (1; 7)	8.7% (4; 48)	13.1% (2; 33)	1.9% (1; 43)	0% (0; 10)	7.7% (6; 83)
	Redwood Memorial Hospital							14.5% (2; 6)	6.3% (2; 32)	12.7% (4; 27)	0% (0; 33)	1.8% (1; 33)	8.4% (7; 85)
	Saint Joseph Hospital – Eureka			0% (0; 28)	0% (0; 16)	1.5% (2; 218)	0% (0; 111)	6.1% (7; 120)	2.5% (5; 113)	15.4% (16; 104)	3.5% (6; 111)	0.7% (1; 76)	2.8% (6; 150)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

### Definition of Terms:

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
Imperial	El Centro Regional Medical Center				11.9% (1; 11)			9.0% (10; 61)	4.7% (9; 170)	11.7% (13; 110)	2.2% (4; 153)	0% (0; 57)	5.6% (16; 218)
	Pioneers Memorial Hospital				18.4% (1; 8)			9.9% (6; 25)	2.6% (4; 145)	9.9% (8; 81)	4.2% (5; 96)	5.0% (1; 31)	4.1% (11; 222)
Inyo	Northern Inyo Hospital							24.0% (1; 5)	0% (0; 29)	13.5% (1; 14)	14.4% (2; 17)	8.5% (2; 24)	<b>22.3%</b> (8; 50) Worse
Kern	Bakersfield Heart Hospital			0% (0; 13)		2.0% (12; 581)	0% (0; 25)	<b>4.8%</b> (25; 485) Better	2.1% (8; 292)	0% (0; 34)	<b>0.5%</b> (1; 78) Better		1.8% (3; 123)
	Bakersfield Memorial Hospital			0% (0; 30)	9.5% (15; 74)	0.8% (4; 401)	0% (0; 82)	6.0% (20; 294)	<b>1.5%</b> (7; 378) Better	7.9% (26; 285)	2.1% (7; 219)	0.8% (1; 140)	<b>2.1%</b> (11; 444) Better
	Delano Regional Medical Center ( <a href="#">View Comment Letter</a> )							7.6% (1; 7)	3.1% (4; 89)	6.0% (2; 30)	5.1% (3; 53)	0% (0; 12)	<b>7.6%</b> (23; 189) Worse
	Good Samaritan Hospital – Bakersfield								0% (0; 54)		0% (0; 11)		2.5% (1; 110)
	Kern Medical Center		0% (0; 4)		6.4% (2; 31)			10.5% (3; 19)	5.9% (4; 95)	6.1% (4; 56)	4.5% (3; 90)	8.3% (1; 17)	2.0% (2; 104)
	Kern Valley Healthcare District							4.9% (1; 8)	1.6% (1; 44)	0% (0; 9)	0% (0; 8)		1.5% (1; 113)
	Mercy Hospital – Bakersfield			25.8% (1; 9)	7.1% (7; 55)		0% (0; 17)	5.7% (2; 19)	1.5% (3; 151)	8.9% (21; 146)	1.7% (4; 234)	<b>0%</b> (0; 96) Better	3.3% (16; 529)
	Ridgecrest Regional Hospital							10.1% (2; 10)	2.7% (2; 101)	14.4% (5; 41)	<b>15.4%</b> (5; 54) Worse		<b>21.8%</b> (11; 101) Worse

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	San Joaquin Community Hospital			0% (0; 3)	<b>10.3%</b> (22; 127) Worse	1.2% (3; 294)	0% (0; 30)	4.8% (10; 230)	2.9% (15; 476)	9.3% (55; 495)	2.3% (7; 298)	1.1% (1; 96)	4.4% (17; 547)
	Tehachapi Hospital								0% (0; 8)				0% (0; 16)
Kings	Central Valley General Hospital							0% (0; 3)	0% (0; 81)	10.3% (2; 27)	0% (0; 32)		2.0% (1; 100)
	Corcoran District Hospital								0% (0; 5)		0% (0; 4)		0% (0; 19)
	Hanford Community Medical Center							10.0% (9; 51)	3.8% (11; 299)	11.7% (8; 77)	<b>5.8%</b> (7; 149) Worse	4.5% (5; 75)	<b>8.8%</b> (25; 381) Worse
Lake	Saint Helena Hospital – Clearlake **								4.7% (1; 45)	20.7% (3; 7)	6.8% (3; 60)	0% (0; 7)	10.3% (8; 117)
	Sutter Lakeside Hospital							14.3% (2; 10)	2.9% (1; 35)	10.6% (4; 54)	2.2% (2; 37)	0% (0; 28)	3.9% (2; 57)
Lassen	Banner Lassen Medical Center								23.6% (2; 19)	7.4% (1; 7)	0% (0; 7)	0% (0; 5)	7.2% (3; 77)
Los Angeles	Alhambra Hospital						0% (0; 3)	7.5% (9; 46)	<b>1.4%</b> (4; 162) Better	<b>5.0%</b> (5; 85) Better	0% (0; 87)	0% (0; 30)	<b>1.3%</b> (5; 177) Better
	Antelope Valley Hospital			0% (0; 3)	6.8% (10; 72)	2.2% (9; 333)	0% (0; 21)	7.1% (27; 369)	2.9% (14; 530)	<b>7.2%</b> (24; 282) Better	3.7% (13; 290)	2.1% (4; 182)	<b>7.8%</b> (60; 657) Worse
	Bellflower Medical Center							8.9% (1; 6)	0% (0; 66)	0% (0; 24)	1.0% (1; 69)	0% (0; 11)	2.8% (5; 124)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

### Definition of Terms:

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurysm Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Beverly Hospital				16.7% (4; 16)	2.0% (4; 141)	0% (0; 19)	4.9% (9; 155)	3.6% (15; 350)	9.7% (13; 155)	3.0% (4; 128)	1.3% (1; 55)	3.1% (12; 284)
	Brotman Medical Center			0% (0; 9)	6.1% (1; 28)	1.7% (2; 62)	0% (0; 11)	5.7% (9; 101)	<b>1.0%</b> (3; 262) <b>Better</b>	10.9% (12; 93)	2.3% (3; 106)	0% (0; 44)	3.9% (13; 220)
	California Hospital Medical Center – Los Angeles				6.1% (4; 17)			8.6% (8; 58)	<b>1.3%</b> (6; 548) <b>Better</b>	8.3% (28; 223)	1.1% (3; 208)	2.9% (1; 34)	4.8% (9; 195)
	Catalina Island Medical Center												0% (0; 3)
	Cedars Sinai Medical Center	8.7% (1; 19)	7.1% (2; 35)	0% (0; 34)	<b>3.9%</b> (23; 601) <b>Better</b>	1.1% (9; 712)	0% (0; 93)	6.8% (33; 464)	3.1% (54; 1,169)	<b>8.4%</b> (69; 611) <b>Better</b>	1.2% (7; 468)	1.5% (5; 210)	3.4% (31; 790)
	Centinela Hospital Medical Center ( <a href="#">View Comment Letter</a> )			3.2% (1; 19)	<b>11.2%</b> (24; 46) <b>Worse</b>	2.6% (6; 263)	0% (0; 6)	5.4% (24; 320)	<b>1.0%</b> (17; 1,311) <b>Better</b>	<b>6.3%</b> (59; 523) <b>Better</b>	1.7% (12; 392)	<b>0.5%</b> (1; 69) <b>Better</b>	<b>2.4%</b> (15; 379) <b>Better</b>
	Citrus Valley Medical Center – Inter Community Campus			32.3% (1; 5)	8.2% (1; 12)	2.9% (8; 270)	0% (0; 56)	6.2% (20; 347)	2.8% (11; 323)	8.7% (13; 174)	2.6% (3; 123)	0.9% (1; 71)	3.6% (8; 199)
	Citrus Valley Medical Center – Queen of the Valley Campus			14.1% (1; 14)	9.4% (4; 48)		4.3% (1; 29)	4.6% (7; 71)	2.3% (11; 370)	9.0% (24; 209)	2.8% (7; 195)	0% (0; 67)	2.4% (6; 245)
	City of Hope Helford Clinical Research Hospital	13.2% (1; 8)	12.8% (1; 10)		<b>0%</b> (0; 121) <b>Better</b>				0% (0; 4)	22.2% (3; 10)	4.0% (1; 21)		<b>13.7%</b> (13; 121) <b>Worse</b>
	Coast Plaza Doctors Hospital							0% (0; 9)	3.0% (4; 113)	11.6% (2; 34)	2.8% (2; 48)	0% (0; 9)	6.6% (11; 146)
	Community Hospital of Long Beach							<b>12.6%</b> (12; 32) <b>Worse</b>	0% (0; 64)	13.0% (8; 71)	0% (0; 52)	0% (0; 20)	6.0% (13; 147)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Community and Mission Hospital of Huntington Park – Slauson							0% (0; 6)	<b>0%</b> (0; 165) Better	0% (0; 15)	0% (0; 46)	0% (0; 6)	3.6% (2; 109)
	Doctors Hospital of West Covina, Inc.										0% (0; 3)		
	Downey Regional Medical Center			0% (0; 7)	5.1% (3; 74)	0% (0; 235)	0% (0; 10)	5.4% (16; 224)	<b>1.9%</b> (14; 561) Better	8.1% (22; 241)	1.3% (5; 286)	0% (0; 105)	3.0% (14; 338)
	East Los Angeles Doctors Hospital							8.4% (8; 47)	0.8% (1; 101)	6.5% (2; 41)	0% (0; 41)	3.7% (2; 14)	3.6% (5; 117)
	East Valley Hospital Medical Center							5.0% (2; 21)	0% (0; 35)	0% (0; 16)	2.9% (1; 15)	15.2% (1; 4)	<b>0.7%</b> (1; 83) Better
	Encino Hospital Medical Center							2.8% (2; 21)	1.7% (4; 84)	5.5% (4; 33)	<b>0%</b> (0; 59) Better	2.2% (2; 24)	2.8% (3; 56)
	Foothill Presbyterian Hospital – Johnston Memorial				9.7% (1; 8)		0% (0; 6)	7.9% (4; 24)	2.5% (7; 186)	<b>6.6%</b> (10; 121) Better	0% (0; 101)	<b>0%</b> (0; 58) Better	<b>1.8%</b> (4; 167) Better
	Garfield Medical Center			0% (0; 6)	10.6% (8; 40)	2.7% (6; 188)	2.9% (1; 24)	6.9% (12; 158)	2.5% (14; 400)	7.6% (18; 246)	3.4% (8; 195)	3.4% (3; 61)	<b>3.0%</b> (18; 341) Better
	Glendale Adventist Medical Center – Wilson Terrace		0% (0; 3)	0% (0; 13)	6.1% (16; 143)	2.0% (11; 589)	0% (0; 15)	5.0% (18; 304)	<b>2.1%</b> (20; 479) Better	<b>8.2%</b> (40; 345) Better	3.1% (11; 261)	2.7% (4; 122)	3.4% (26; 437)
	Glendale Memorial Hospital and Medical Center	63.6% (1; 3)		0% (0; 12)	4.8% (2; 20)	1.7% (6; 310)	0% (0; 25)	<b>3.9%</b> (11; 264) Better	<b>1.9%</b> (18; 565) Better	<b>5.7%</b> (15; 201) Better	1.7% (4; 183)	2.1% (4; 125)	3.6% (18; 296)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Good Samaritan Hospital – Los Angeles			0% (0; 9)	5.9% (4; 45)	<b>3.3%</b> (15; 418) Worse	6.7% (1; 10)	9.2% (25; 305)	3.1% (21; 684)	9.6% (20; 185)	2.3% (5; 222)	1.2% (1; 82)	4.8% (19; 263)
	Greater El Monte Community Hospital							3.8% (3; 23)	2.4% (2; 75)	7.7% (4; 42)	0% (0; 43)	0% (0; 3)	<b>1.9%</b> (4; 118) Better
	Harbor – UCLA Medical Center		0% (0; 6)	0% (0; 11)	4.0% (3; 100)	4.9% (5; 300)	0% (0; 10)	11.5% (13; 239)	<b>0.6%</b> (2; 549) Better	9.8% (23; 264)	2.4% (5; 191)	3.4% (2; 34)	4.2% (10; 378)
	Henry Mayo Newhall Memorial Hospital				11.7% (5; 32)		0% (0; 11)	<b>3.6%</b> (7; 95) Better	3.2% (14; 259)	9.3% (20; 183)	0.5% (1; 159)	1.4% (2; 85)	4.8% (15; 275)
	Hollywood Community Hospital							19.9% (1; 7)	0% (0; 67)	0% (0; 13)	0% (0; 37)	0% (0; 4)	4.6% (6; 129)
	Hollywood Presbyterian Medical Center		0% (0; 4)		2.0% (1; 20)		0% (0; 4)	7.2% (16; 111)	2.7% (16; 433)	7.6% (17; 137)	3.6% (9; 191)	0% (0; 74)	5.9% (19; 253)
	Huntington Memorial Hospital		0% (0; 5)	0% (0; 15)	9.6% (12; 160)	1.9% (7; 321)	0% (0; 40)	4.6% (14; 274)	<b>1.8%</b> (13; 586) Better	9.4% (35; 360)	3.0% (10; 323)	1.0% (2; 177)	<b>2.2%</b> (9; 407) Better
	Kaiser Foundation Hospital – Baldwin Park			0% (0; 19)			0% (0; 51)	6.3% (10; 115)	<b>1.2%</b> (7; 483) Better	<b>6.0%</b> (11; 217) Better	1.5% (5; 302)	1.3% (2; 92)	3.7% (17; 453)
	Kaiser Foundation Hospital – Bellflower	0% (0; 4)		0% (0; 23)			0% (0; 32)	4.3% (5; 87)	<b>1.4%</b> (5; 340) Better	6.1% (7; 152)	0.7% (2; 255)	3.0% (3; 65)	<b>2.1%</b> (6; 306) Better
	Kaiser Foundation Hospital – Downey			0% (0; 8)			0% (0; 15)	2.0% (1; 43)	2.1% (3; 118)	11.7% (10; 85)	2.4% (2; 105)	0% (0; 33)	1.1% (1; 109)
	Kaiser Foundation Hospital – Harbor City	0% (0; 6)	25.5% (1; 8)	0% (0; 15)			0% (0; 54)	5.7% (9; 89)	<b>1.4%</b> (8; 499) Better	8.2% (15; 208)	1.1% (4; 336)	2.9% (3; 66)	<b>2.5%</b> (11; 379) Better

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

### Definition of Terms:

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Kaiser Foundation Hospital – Panorama City	0% (0; 3)	0% (0; 8)	0% (0; 6)		0% (0; 28)	0% (0; 30)	5.7% (7; 82)	<b>1.2%</b> (6; 422) Better	<b>6.6%</b> (14; 233) Better	1.6% (7; 343)	2.3% (4; 129)	3.4% (20; 548)
	Kaiser Foundation Hospital – Sunset		0% (0; 13)	3.5% (1; 31)	5.5% (16; 406)	1.2% (9; 852)	0% (0; 11)	<b>5.2%</b> (39; 1,009) Better	2.8% (20; 560)	<b>7.5%</b> (45; 471) Better	<b>0.6%</b> (2; 245) Better	1.6% (1; 62)	3.7% (13; 325)
	Kaiser Foundation Hospital – West Los Angeles	0% (0; 4)		0% (0; 7)			0% (0; 16)	5.6% (10; 116)	<b>0.8%</b> (4; 475) Better	8.5% (23; 286)	2.3% (6; 338)	0% (0; 62)	4.6% (24; 358)
	Kaiser Foundation Hospital – Woodland Hills			6.9% (1; 11)	3.8% (4; 75)		1.0% (1; 92)	7.3% (12; 106)	2.6% (24; 426)	<b>6.3%</b> (25; 285) Better	1.9% (7; 270)	2.0% (3; 145)	<b>1.7%</b> (8; 349) Better
	Lakewood Regional Medical Center				10.2% (4; 33)	1.6% (4; 347)	7.2% (1; 22)	4.4% (9; 217)	2.9% (12; 447)	10.5% (21; 204)	3.2% (6; 199)	2.6% (2; 87)	4.0% (14; 241)
	Lancaster Community Hospital			<b>21.1%</b> (2; 21) Worse	6.6% (2; 16)	0.8% (1; 209)	0% (0; 8)	5.3% (7; 152)	1.4% (4; 286)	7.2% (10; 123)	3.4% (5; 134)	1.7% (1; 55)	3.7% (10; 266)
	Long Beach Memorial Medical Center			0% (0; 34)	4.5% (6; 112)	2.1% (11; 599)	1.0% (1; 104)	8.7% (26; 371)	<b>1.7%</b> (12; 706) Better	<b>7.2%</b> (35; 429) Better	2.8% (12; 352)	<b>0%</b> (0; 134) Better	3.7% (26; 557)
	Los Angeles Community Hospital							<b>19.5%</b> (5; 25) Worse	0.9% (1; 123)	5.2% (3; 83)	1.6% (1; 84)		3.8% (8; 161)
	Los Angeles County/University of Southern California Medical Center	0% (0; 3)	18.9% (1; 9)		10.9% (11; 144)	<b>4.3%</b> (9; 260) Worse	0% (0; 14)	<b>11.9%</b> (15; 194) Worse	3.0% (12; 717)	12.1% (30; 225)	1.4% (2; 262)	0% (0; 34)	6.2% (22; 539)
	Los Angeles Metropolitan Medical Center								1.1% (1; 131)	0% (0; 13)	1.9% (1; 40)		3.7% (8; 157)
	Marina Del Rey Hospital						0% (0; 8)	4.0% (3; 46)	1.9% (5; 195)	5.7% (4; 53)	3.9% (4; 90)	0% (0; 40)	5.4% (7; 110)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Memorial Hospital of Gardena				13.0% (5; 11)			10.5% (6; 33)	<b>0.9%</b> (2; 249) <b>Better</b>	6.9% (12; 147)	3.6% (9; 134)	0% (0; 17)	2.6% (4; 166)
	Methodist Hospital of Southern California			4.3% (1; 9)	2.4% (1; 33)	1.9% (8; 316)	0% (0; 28)	6.0% (22; 299)	2.6% (22; 606)	<b>7.1%</b> (27; 295) <b>Better</b>	2.3% (7; 234)	2.1% (4; 199)	3.9% (21; 448)
	Mission Community Hospital – Panorama Campus							12.4% (6; 15)	<b>0.5%</b> (1; 124) <b>Better</b>	<b>3.1%</b> (2; 49) <b>Better</b>	0.7% (1; 80)	0% (0; 25)	<b>1.5%</b> (3; 130) <b>Better</b>
	Monrovia Medical Center								<b>14.4%</b> (3; 3) <b>Worse</b>				4.8% (5; 32)
	Monterey Park Hospital							10.0% (7; 36)	<b>0.9%</b> (2; 147) <b>Better</b>	3.4% (1; 53)	0.8% (1; 108)	0% (0; 29)	4.0% (4; 97)
	Motion Picture and Television Hospital								6.4% (1; 19)	18.9% (1; 10)	0% (0; 13)		4.9% (1; 28)
	Northridge Hospital Medical Center			0% (0; 10)	9.3% (7; 46)	2.4% (8; 354)	0% (0; 23)	7.7% (21; 221)	2.0% (11; 378)	10.8% (40; 246)	1.5% (4; 216)	1.6% (3; 114)	<b>2.8%</b> (13; 346) <b>Better</b>
	Norwalk Community Hospital							12.2% (2; 18)	1.8% (2; 109)	7.5% (2; 35)	0% (0; 55)	0% (0; 10)	2.6% (3; 76)
	Olive View – UCLA Medical Center			0% (0; 3)			<b>52.0%</b> (1; 11) <b>Worse</b>	7.0% (3; 39)	2.7% (6; 326)	10.9% (10; 114)	0% (0; 134)		2.5% (6; 388)
	Olympia Medical Center				0% (0; 4)			<b>3.7%</b> (9; 114) <b>Better</b>	3.2% (12; 280)	5.5% (5; 95)	3.4% (10; 153)	3.4% (3; 64)	<b>1.9%</b> (10; 295) <b>Better</b>

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Pacific Alliance Medical Center, Inc.				0% (0; 5)			4.1% (1; 17)	2.3% (9; 209)	<b>1.5%</b> (1; 74) <b>Better</b>	2.7% (3; 81)	3.9% (1; 12)	3.8% (6; 110)
	Pacific Hospital of Long Beach							9.9% (1; 7)	0% (0; 94)	6.2% (1; 21)	3.1% (3; 74)	0% (0; 13)	5.2% (10; 122)
	Pacifica Hospital of the Valley							6.8% (1; 8)	3.5% (3; 94)	<b>15.6%</b> (13; 58) <b>Worse</b>	4.1% (2; 61)	3.7% (1; 16)	11.1% (6; 77)
	Pomona Valley Hospital Medical Center		0% (0; 3)	0% (0; 19)	3.3% (4; 80)	1.7% (6; 232)	0% (0; 16)	5.1% (12; 241)	2.2% (17; 578)	<b>6.1%</b> (27; 287) <b>Better</b>	1.1% (4; 311)	0% (0; 128)	2.7% (9; 261)
	Presbyterian Intercommunity Hospital	0% (0; 3)	63.0% (1; 4)	12.9% (2; 14)	<b>0%</b> (0; 56) <b>Better</b>	1.0% (4; 299)	0% (0; 48)	<b>3.9%</b> (28; 480) <b>Better</b>	<b>1.4%</b> (9; 466) <b>Better</b>	<b>6.3%</b> (21; 275) <b>Better</b>	<b>0.6%</b> (2; 300) <b>Better</b>	1.4% (2; 133)	<b>2.1%</b> (10; 422) <b>Better</b>
	Providence Holy Cross Medical Center	0% (0; 4)	0% (0; 3)	0% (0; 9)	5.5% (4; 44)	2.2% (9; 343)	0% (0; 69)	7.4% (16; 283)	2.4% (19; 503)	<b>7.0%</b> (19; 226) <b>Better</b>	1.3% (5; 254)	2.6% (2; 82)	3.1% (18; 369)
	Providence Little Company of Mary Medical Center – San Pedro**				0% (0; 9)		0% (0; 10)	4.2% (2; 28)	3.7% (7; 163)	6.1% (6; 92)	2.5% (3; 104)	1.0% (1; 71)	5.1% (12; 183)
	Providence Little Company of Mary Medical Center – Torrance**		0% (0; 3)	0% (0; 7)	16.0% (3; 26)	2.5% (14; 491)	0% (0; 23)	8.1% (23; 267)	3.1% (24; 602)	9.0% (37; 303)	1.5% (6; 352)	0.7% (1; 143)	5.0% (26; 450)
	Providence Saint Joseph Medical Center			0% (0; 17)	3.8% (4; 89)	2.0% (11; 345)	0% (0; 62)	5.2% (14; 272)	3.3% (36; 619)	11.6% (46; 338)	2.4% (8; 275)	1.0% (2; 180)	5.5% (22; 301)
	Providence Tarzana Medical Center			0% (0; 13)	16.5% (1; 11)	3.2% (6; 220)	2.3% (2; 107)	10.4% (16; 168)	3.5% (22; 334)	9.3% (15; 133)	2.6% (4; 173)	2.5% (5; 130)	3.7% (18; 327)
	Ronald Reagan UCLA Medical Center	0% (0; 22)	4.6% (2; 63)	0% (0; 75)	5.3% (34; 600)	2.6% (5; 152)	0% (0; 42)	<b>3.0%</b> (6; 175) <b>Better</b>	4.5% (24; 489)	<b>8.3%</b> (56; 398) <b>Better</b>	2.4% (5; 127)	4.0% (2; 39)	6.5% (18; 207)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Saint Francis Medical Center			0% (0; 4)	11.9% (7; 34)	1.1% (2; 139)	0% (0; 20)	6.5% (16; 158)	2.2% (14; 614)	8.9% (43; 264)	<b>0%</b> <b>(0; 202)</b> <b>Better</b>	0% (0; 26)	3.5% (10; 253)
	Saint John's Health Center	0% (0; 6)	0% (0; 10)	0% (0; 17)	2.8% (2; 83)	0.6% (1; 289)	0% (0; 60)	5.2% (13; 192)	<b>1.6%</b> <b>(8; 333)</b> <b>Better</b>	7.4% (15; 181)	1.1% (3; 219)	1.8% (3; 154)	3.3% (17; 338)
	Saint Mary Medical Center	<b>33.6%</b> <b>(2; 5)</b> <b>Worse</b>		35.6% (1; 4)	5.6% (2; 23)	4.0% (5; 148)	0% (0; 24)	6.7% (11; 146)	1.9% (7; 286)	8.9% (25; 185)	<b>0%</b> <b>(0; 176)</b> <b>Better</b>	0.8% (1; 71)	3.2% (13; 315)
	Saint Vincent Medical Center		0% (0; 12)	8.4% (1; 14)	3.5% (3; 259)	1.2% (2; 197)	4.0% (1; 13)	9.3% (12; 114)	2.1% (10; 413)	<b>4.5%</b> <b>(6; 138)</b> <b>Better</b>	1.4% (2; 145)	1.2% (1; 47)	3.3% (12; 331)
	San Dimas Community Hospital ( <a href="#">View Comment Letter</a> )						0% (0; 3)	7.4% (8; 47)	2.9% (4; 108)	<b>4.9%</b> <b>(4; 77)</b> <b>Better</b>	0.7% (1; 100)	3.5% (1; 35)	3.9% (4; 77)
	San Gabriel Valley Medical Center				3.4% (1; 18)			5.0% (5; 38)	1.8% (9; 270)	<b>6.6%</b> <b>(13; 171)</b> <b>Better</b>	0.8% (2; 124)	1.4% (1; 64)	4.3% (28; 307)
	Santa Monica – UCLA Medical Center and Orthopedic Hospital				17.0% (2; 15)	2.3% (6; 130)	0% (0; 6)	9.2% (22; 159)	<b>1.3%</b> <b>(8; 367)</b> <b>Better</b>	<b>6.3%</b> <b>(10; 166)</b> <b>Better</b>	1.4% (4; 232)	1.1% (2; 140)	<b>2.5%</b> <b>(12; 354)</b> <b>Better</b>
	Sherman Oaks Hospital						0% (0; 3)	<b>2.0%</b> <b>(7; 111)</b> <b>Better</b>	<b>0.7%</b> <b>(3; 157)</b> <b>Better</b>	<b>5.4%</b> <b>(6; 56)</b> <b>Better</b>	1.1% (2; 70)	0% (0; 42)	<b>0%</b> <b>(0; 58)</b> <b>Better</b>
	Silver Lake Medical Center – Downtown Campus							0% (0; 7)	0% (0; 51)	0% (0; 5)	3.2% (1; 39)		8.7% (4; 47)
	Temple Community Hospital							5.1% (2; 21)	3.5% (6; 131)	0% (0; 27)	2.2% (1; 52)		8.8% (6; 52)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

### Definition of Terms:

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Torrance Memorial Medical Center ( <a href="#">View Comment Letter</a> )		0% (0; 3)	0% (0; 32)	5.5% (3; 46)	1.1% (6; 622)	4.6% (2; 87)	6.6% (31; 388)	3.3% (27; 646)	8.6% (31; 408)	1.8% (8; 422)	1.1% (2; 201)	3.9% (24; 506)
	Tri-City Regional Medical Center – Hawaiian Gardens						0% (0; 4)	7.7% (1; 4)	1.2% (2; 120)	34.1% (2; 15)	1.5% (1; 65)	0% (0; 12)	5.2% (7; 132)
	University of Southern California Kenneth Norris, Jr. Cancer Hospital								0% (0; 5)		13.0% (2; 20)		6.4% (1; 20)
	University of Southern California University Hospital	3.4% (1; 42)	9.7% (2; 35)	0% (0; 33)	3.2% (5; 258)	<b>7.2%</b> (7; 74) Worse	0% (0; 16)	10.3% (6; 60)	5.7% (10; 119)	8.7% (10; 97)	0% (0; 28)	0% (0; 9)	3.5% (2; 47)
	Valley Presbyterian Hospital			0% (0; 8)	6.7% (3; 16)	2.2% (6; 246)	0% (0; 17)	6.9% (16; 227)	3.3% (18; 377)	<b>13.8%</b> (34; 179) Worse	2.4% (4; 142)	4.6% (4; 78)	5.0% (20; 259)
	Verdugo Hills Hospital				0% (0; 11)		0% (0; 10)	9.7% (5; 37)	<b>0.5%</b> (1; 143) Better	7.6% (9; 87)	2.5% (2; 97)	3.7% (3; 84)	5.0% (11; 218)
	West Hills Hospital and Medical Center				10.4% (4; 22)	1.8% (3; 129)	0% (0; 26)	10.0% (14; 133)	3.6% (12; 208)	10.7% (20; 160)	3.2% (7; 202)	4.1% (5; 102)	<b>2.8%</b> (14; 258) Better
	White Memorial Medical Center				3.6% (1; 21)	1.6% (9; 326)	0% (0; 17)	<b>4.5%</b> (15; 236) Better	2.0% (10; 388)	<b>7.1%</b> (17; 182) Better	1.2% (5; 231)	<b>0%</b> (0; 50) Better	<b>1.4%</b> (6; 297) Better
	Whittier Hospital Medical Center				0% (0; 3)		0% (0; 5)	<b>1.9%</b> (4; 98) Better	1.8% (6; 222)	12.6% (8; 104)	<b>0.6%</b> (2; 202) Better	1.6% (2; 90)	1.4% (2; 182)
Madera	Madera Community Hospital							11.7% (13; 73)	<b>9.0%</b> (15; 131) Worse	<b>31.8%</b> (6; 34) Worse	1.6% (1; 61)	6.4% (1; 13)	4.5% (7; 200)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurysm Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	5.8% (29; 500)	5.2% (38; 732)	1.9% (50; 2,567)	6.7% (938; 13,926)	1.9% (933; 48,689)	0.7% (49; 7,411)	7.1% (3,151; 44,510)	3.3% (2,832; 85,934)	10.4% (5,285; 50,988)	2.3% (1,114; 47,720)	2.4% (571; 23,514)	4.6% (3,703; 79,958)
Marin	Kaiser Foundation Hospital – San Rafael					0% (0; 29)	0% (0; 19)	2.3% (2; 89)	3.2% (10; 203)	12.2% (18; 170)	2.0% (2; 125)	2.7% (5; 98)	6.8% (21; 274)
	Marin General Hospital			0% (0; 23)	8.3% (6; 29)	0.7% (1; 214)	<b>7.3% (2; 51) Worse</b>	4.3% (6; 181)	2.3% (7; 200)	10.5% (25; 182)	1.5% (3; 179)	1.8% (3; 98)	3.5% (8; 187)
	Novato Community Hospital						0% (0; 6)	2.2% (1; 17)	2.1% (2; 69)	10.8% (3; 25)	0% (0; 43)	2.5% (1; 36)	6.9% (5; 68)
Mariposa	John C. Fremont Healthcare District							0% (0; 4)	9.0% (1; 11)	<b>85.7% (3; 5) Worse</b>	0% (0; 4)		0% (0; 20)
Mendocino	Frank R. Howard Memorial Hospital							0% (0; 4)	<b>14.3% (4; 22) Worse</b>	6.8% (1; 17)	0% (0; 18)	3.6% (1; 35)	0% (0; 42)
	Mendocino Coast District Hospital							0% (0; 6)	10.6% (4; 31)	9.2% (2; 25)	0% (0; 20)	0% (0; 9)	0% (0; 57)
	Ukiah Valley Medical Center						0% (0; 6)	9.3% (3; 14)	<b>1.1% (3; 103) Better</b>	7.2% (6; 56)	1.1% (1; 51)	4.6% (3; 36)	<b>1.8% (5; 172) Better</b>
Merced	Memorial Hospital Los Banos							4.0% (1; 8)	0% (0; 39)	4.7% (2; 26)	0% (0; 20)		2.7% (3; 85)
	Mercy Medical Center – Merced							8.5% (13; 82)	3.2% (17; 330)	7.6% (10; 160)	3.5% (12; 228)	2.4% (2; 76)	4.2% (17; 348)
Modoc	Modoc Medical Center								17.8% (2; 12)		0% (0; 3)		0% (0; 25)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.8% (29; 500)	5.2% (38; 732)	1.9% (50; 2,567)	6.7% (938; 13,926)	1.9% (933; 48,689)	0.7% (49; 7,411)	7.1% (3,151; 44,510)	3.3% (2,832; 85,934)	10.4% (5,285; 50,988)	2.3% (1,114; 47,720)	2.4% (571; 23,514)	4.6% (3,703; 79,958)
	Surprise Valley Community Hospital												42.1% (1; 7)
Mono	Mammoth Hospital								0% (0; 5)		0% (0; 6)	0% (0; 17)	0% (0; 13)
Monterey	Community Hospital Monterey Peninsula		0% (0; 3)	0% (0; 4)	9.7% (3; 33)	1.0% (2; 173)	0% (0; 44)	8.0% (16; 181)	2.9% (10; 259)	12.7% (31; 186)	4.0% (10; 234)	3.0% (5; 126)	5.4% (18; 305)
	George L. Mee Memorial Hospital							8.5% (1; 6)	0% (0; 36)	18.1% (2; 5)	7.0% (1; 15)		4.8% (3; 56)
	Natividad Medical Center							15.1% (7; 27)	1.1% (1; 85)	6.1% (1; 33)	0% (0; 90)	0% (0; 10)	4.5% (6; 151)
	Salinas Valley Memorial Hospital	0% (0; 4)		0% (0; 6)	3.6% (2; 41)	1.7% (5; 222)	0% (0; 32)	5.5% (13; 233)	2.8% (10; 310)	9.0% (24; 202)	1.8% (5; 225)	3.3% (4; 101)	3.2% (11; 265)
Napa	N.M. Holderman Memorial Hospital (Veteran's Home of California) <a href="#">(View Comment Letter)</a>												0% (0; 6)
	Queen of the Valley Hospital – Napa			0% (0; 17)		2.8% (7; 190)	0% (0; 37)	5.8% (11; 204)	3.2% (13; 224)	11.5% (11; 98)	1.9% (4; 168)	2.5% (4; 85)	3.3% (10; 245)
	Saint Helena Hospital			0% (0; 17)		1.2% (2; 339)	1.0% (1; 85)	9.0% (6; 71)	3.0% (4; 103)	9.4% (4; 45)	2.3% (3; 68)	3.4% (1; 27)	3.0% (4; 109)
Nevada	Sierra Nevada Memorial Hospital						0% (0; 12)	8.1% (6; 38)	2.5% (6; 160)	10.4% (8; 91)	3.8% (4; 99)	2.5% (4; 113)	3.2% (6; 191)
	Tahoe Forest Hospital								5.4% (1; 26)	0% (0; 9)	0% (0; 15)	0% (0; 33)	4.6% (1; 38)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
Orange	AHMC Anaheim Regional Medical Center**			0% (0; 25)	7.9% (2; 20)	1.6% (5; 337)	1.3% (1; 41)	7.0% (23; 300)	2.0% (11; 480)	<b>5.9%</b> (10; 181) Better	2.5% (6; 225)	0.7% (1; 97)	<b>2.2%</b> (9; 363) Better
	Anaheim General Hospital							0% (0; 3)	0% (0; 19)	0% (0; 5)	0% (0; 20)		1.2% (1; 44)
	Chapman Medical Center								0% (0; 46)	12.6% (2; 12)	0% (0; 34)	0% (0; 9)	3.9% (5; 79)
	Coastal Communities Hospital						0% (0; 3)	14.4% (5; 16)	2.1% (2; 83)	5.4% (1; 23)	2.2% (2; 61)	0% (0; 11)	<b>9.4%</b> (15; 119) Worse
	Fountain Valley Regional Hospital and Medical Center – Euclid	0% (0; 4)	0% (0; 3)	0% (0; 10)	4.6% (5; 67)	0.9% (3; 274)	0% (0; 26)	5.3% (13; 290)	<b>1.5%</b> (8; 489) Better	9.3% (28; 310)	<b>5.2%</b> (16; 283) Worse	0.8% (1; 99)	3.8% (16; 363)
	Garden Grove Hospital and Medical Center				0% (0; 12)		0% (0; 3)	<b>3.5%</b> (7; 78) Better	2.0% (3; 110)	<b>4.5%</b> (6; 102) Better	2.1% (2; 101)	3.5% (1; 35)	3.5% (5; 114)
	Hoag Memorial Hospital Presbyterian	0% (0; 3)		0% (0; 28)	6.2% (20; 300)	1.6% (5; 505)	0% (0; 89)	7.9% (19; 260)	2.7% (17; 492)	8.4% (46; 508)	3.5% (17; 387)	1.6% (5; 298)	3.9% (31; 606)
	Huntington Beach Hospital							5.5% (10; 49)	1.3% (3; 119)	3.4% (1; 40)	2.0% (4; 69)	0% (0; 24)	<b>1.5%</b> (4; 117) Better
	Irvine Medical Center					0% (0; 5)			15.5% (1; 5)				0% (0; 9)
	Kaiser Foundation Hospital – Anaheim		14.6% (1; 8)	0% (0; 17)	5.1% (7; 194)		0% (0; 49)	<b>3.7%</b> (11; 168) Better	<b>1.7%</b> (12; 484) Better	<b>6.8%</b> (36; 471) Better	1.2% (5; 383)	1.3% (3; 193)	<b>2.8%</b> (21; 602) Better
	La Palma Intercommunity Hospital ( <a href="#">View Comment Letter</a> )							3.7% (5; 62)	1.5% (3; 163)	2.8% (1; 64)	0% (0; 78)	0% (0; 31)	2.4% (2; 77)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurysm Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Los Alamitos Medical Center			0% (0; 18)	6.7% (3; 29)	3.7% (3; 58)	1.8% (1; 75)	11.1% (16; 125)	3.2% (13; 371)	8.2% (24; 270)	2.2% (4; 163)	1.5% (3; 135)	7.0% (22; 329)
	Mission Hospital Laguna Beach				0% (0; 5)		<b>27.2%</b> (1; 3) Worse	7.0% (2; 10)	0% (0; 62)	5.0% (1; 20)	5.4% (3; 44)	2.6% (1; 28)	4.7% (3; 59)
	Mission Hospital Regional Medical Center			0% (0; 15)	7.0% (13; 117)	1.5% (6; 506)	0% (0; 15)	7.0% (17; 274)	3.7% (20; 326)	10.5% (48; 350)	0.8% (2; 230)	1.9% (4; 159)	5.0% (14; 308)
	Orange Coast Memorial Medical Center ( <a href="#">View Comment Letter</a> )		0% (0; 6)		2.6% (1; 29)	16.4% (1; 19)	0% (0; 20)	4.2% (9; 121)	2.0% (11; 378)	<b>3.6%</b> (5; 165) Better	2.3% (5; 217)	0.8% (1; 132)	5.4% (17; 255)
	Placentia Linda Hospital						0% (0; 3)	7.2% (5; 31)	2.7% (4; 146)	6.0% (3; 71)	0% (0; 111)	0% (0; 72)	<b>1.6%</b> (3; 174) Better
	Saddleback Memorial Medical Center			0% (0; 21)	3.2% (3; 69)	1.7% (9; 327)	0% (0; 34)	<b>5.0%</b> (26; 349) Better	<b>1.7%</b> (18; 585) Better	<b>6.5%</b> (26; 332) Better	1.4% (5; 319)	1.7% (10; 331)	3.2% (20; 463)
	Saint Joseph Hospital – Orange	0% (0; 9)	13.6% (1; 10)	0% (0; 28)	<b>3.0%</b> (8; 161) Better	1.7% (6; 269)	0% (0; 91)	6.3% (18; 234)	2.1% (15; 441)	<b>5.0%</b> (22; 323) Better	1.6% (7; 287)	1.4% (3; 174)	4.4% (16; 286)
	Saint Jude Medical Center			0% (0; 19)	8.7% (8; 69)	1.5% (8; 464)	1.5% (1; 73)	6.2% (28; 421)	2.8% (20; 470)	9.1% (43; 405)	1.7% (7; 288)	2.6% (7; 199)	3.6% (19; 412)
	UC Irvine Medical Center	6.6% (1; 19)	5.1% (1; 25)	13.3% (2; 17)	6.1% (19; 209)	2.3% (4; 165)	4.5% (1; 33)	8.5% (9; 123)	2.3% (5; 220)	<b>8.6%</b> (59; 333) Better	1.9% (4; 151)	1.7% (1; 36)	3.2% (7; 209)
	West Anaheim Medical Center					1.4% (3; 124)	0% (0; 3)	4.7% (9; 130)	<b>0.6%</b> (3; 232) Better	<b>0%</b> (0; 82) Better	2.3% (6; 136)	0% (0; 46)	<b>1.6%</b> (8; 233) Better

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

### Definition of Terms:

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	5.8% (29; 500)	5.2% (38; 732)	1.9% (50; 2,567)	6.7% (938; 13,926)	1.9% (933; 48,689)	0.7% (49; 7,411)	7.1% (3,151; 44,510)	3.3% (2,832; 85,934)	10.4% (5,285; 50,988)	2.3% (1,114; 47,720)	2.4% (571; 23,514)	4.6% (3,703; 79,958)
	Western Medical Center – Anaheim					2.1% (2; 130)	0% (0; 5)	3.3% (2; 112)	1.7% (1; 57)	0% (0; 17)	1.8% (1; 53)	0% (0; 14)	3.2% (2; 65)
	Western Medical Center – Santa Ana				6.9% (6; 55)	0.9% (5; 528)	0% (0; 15)	3.4% (10; 382) <b>Better</b>	5.1% (12; 200)	9.2% (31; 198)	0.8% (1; 102)	5.1% (3; 44)	2.3% (4; 149)
Placer	Sutter Auburn Faith Hospital			0% (0; 8)			0% (0; 19)	5.9% (2; 14)	3.8% (8; 102)	7.3% (6; 71)	2.6% (2; 46)	1.0% (1; 84)	4.5% (7; 151)
	Sutter Roseville Medical Center	0% (0; 3)	0% (0; 3)	0% (0; 13)	7.3% (9; 88)	2.4% (5; 148)	3.7% (1; 52)	7.1% (20; 193)	2.7% (14; 315)	9.8% (41; 318)	1.8% (6; 271)	3.5% (10; 186)	2.1% (10; 372) <b>Better</b>
Plumas	Eastern Plumas Hospital – Portola Campus							30.3% (1; 3)	16.4% (5; 24) <b>Worse</b>		0% (0; 6)		5.0% (1; 30)
	Plumas District Hospital								0% (0; 14)	0% (0; 7)	0% (0; 12)		0% (0; 34)
	Seneca Healthcare District								0% (0; 15)	0% (0; 4)	0% (0; 3)		12.6% (2; 24)
Riverside	Corona Regional Medical Center – Main						0% (0; 11)	6.0% (6; 64)	3.2% (12; 270)	7.6% (5; 105)	3.2% (5; 147)	2.0% (2; 83)	6.3% (23; 307)
	Desert Regional Medical Center	15.3% (1; 4)		0% (0; 18)	11.2% (9; 50)	1.7% (12; 560)	0% (0; 56)	6.3% (17; 361)	4.3% (16; 336)	9.6% (63; 408)	2.4% (9; 291)	3.6% (7; 166)	7.5% (28; 346) <b>Worse</b>
	Eisenhower Medical Center			5.9% (1; 20)	2.6% (2; 74)	2.4% (15; 670)	0% (0; 99)	7.9% (34; 447)	3.0% (23; 612)	7.5% (34; 473) <b>Better</b>	1.6% (7; 429)	1.9% (5; 251)	3.3% (23; 595)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Hemet Valley Medical Center ( <a href="#">View Comment Letter</a> )			0% (0; 5)			<b>6.2%</b> (2; 22) Worse	7.7% (32; 196)	3.5% (19; 415)	10.1% (18; 221)	1.5% (4; 264)	1.5% (3; 186)	5.4% (25; 340)
	John F. Kennedy Memorial Hospital					0% (0; 12)	0% (0; 12)	11.9% (12; 69)	1.2% (2; 184)	5.5% (2; 52)	3.2% (3; 90)	3.5% (2; 62)	2.4% (3; 181)
	Kaiser Foundation Hospital – Moreno Valley						0% (0; 9)	13.3% (3; 13)	5.1% (5; 108)	0% (0; 41)	1.7% (1; 58)	12.9% (1; 5)	<b>1.3%</b> (2; 139) Better
	Kaiser Foundation Hospital – Riverside	0% (0; 9)	0% (0; 14)	0% (0; 10)			0% (0; 40)	3.7% (6; 109)	3.4% (17; 387)	7.5% (12; 218)	1.2% (3; 268)	0% (0; 98)	4.1% (19; 396)
	Menifee Valley Medical Center ( <a href="#">View Comment Letter</a> )						0% (0; 11)	6.3% (12; 76)	2.3% (5; 197)	5.5% (3; 75)	3.9% (4; 119)	<b>7.3%</b> (6; 84) Worse	4.9% (8; 148)
	Palo Verde Hospital							24.0% (2; 7)	4.6% (1; 24)	0% (0; 18)	0% (0; 41)		11.4% (4; 81)
	Parkview Community Hospital Medical Center						0% (0; 8)	8.7% (6; 40)	3.4% (9; 182)	7.7% (7; 83)	0.7% (1; 134)	0% (0; 61)	4.2% (16; 299)
	Riverside Community Hospital			0% (0; 20)	6.2% (4; 64)	2.2% (30; 848)	0% (0; 38)	5.9% (34; 690)	2.7% (19; 612)	9.1% (33; 303)	2.4% (7; 289)	<b>0%</b> (0; 150) Better	5.6% (21; 357)
	Riverside County Regional Medical Center			0% (0; 3)	4.4% (1; 38)		0% (0; 9)	7.2% (4; 63)	1.7% (4; 388)	7.6% (17; 168)	2.5% (4; 144)	3.4% (1; 43)	2.8% (8; 325)
	San Geronio Memorial Hospital							5.5% (8; 68)	2.9% (6; 167)	<b>1.9%</b> (1; 73) Better	0.8% (1; 84)	0% (0; 30)	5.5% (11; 193)
	Southwest Healthcare System – Murrieta			0% (0; 31)	15.7% (6; 60)		0% (0; 94)	7.4% (18; 128)	2.3% (11; 498)	10.1% (29; 295)	0.9% (2; 250)	1.6% (2; 159)	4.0% (18; 481)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurysm Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
Sacramento	Kaiser Foundation Hospital – Sacramento/Roseville-Morse	0% (0; 12)	16.3% (1; 7)	<b>21.0%</b> (4; 33) Worse	<b>4.4%</b> (15; 293) Better	0% (0; 28)	0% (0; 94)	6.4% (27; 307)	<b>4.5%</b> (50; 931) Worse	10.1% (90; 786)	<b>1.0%</b> (5; 529) Better	2.2% (9; 385)	5.2% (40; 771)
	Kaiser Foundation Hospital – South Sacramento		0% (0; 4)	13.9% (2; 25)	0% (0; 6)		0% (0; 37)	6.8% (13; 140)	<b>1.5%</b> (6; 358) Better	<b>6.6%</b> (13; 239) Better	0.7% (1; 203)	5.7% (5; 106)	4.2% (12; 294)
	Mercy General Hospital			0% (0; 24)	8.7% (12; 101)	1.9% (24; 1,631)	0.6% (1; 105)	5.4% (37; 1,147)	1.9% (11; 458)	10.7% (35; 294)	1.3% (3; 205)	0.8% (1; 115)	3.3% (12; 299)
	Mercy Hospital – Folsom				10.3% (1; 5)			10.9% (4; 11)	1.2% (2; 96)	<b>4.3%</b> (3; 83) Better	0% (0; 90)	3.5% (2; 65)	4.0% (6; 144)
	Mercy San Juan Hospital			4.8% (1; 28)	6.6% (6; 61)	1.0% (2; 233)	0% (0; 54)	6.1% (22; 315)	2.9% (28; 509)	12.2% (71; 427)	3.3% (14; 299)	1.8% (4; 204)	3.7% (17; 406)
	Methodist Hospital of Sacramento	0% (0; 3)					0% (0; 4)	7.4% (7; 34)	2.4% (9; 264)	8.7% (19; 163)	4.2% (10; 171)	1.2% (1; 86)	5.0% (13; 251)
	Sutter General Hospital	8.8% (1; 9)	0% (0; 8)	0% (0; 10)	4.6% (6; 164)		0% (0; 34)	11.9% (6; 21)	1.8% (7; 284)	<b>6.9%</b> (20; 251) Better	1.4% (4; 211)	3.2% (7; 176)	<b>1.3%</b> (5; 399) Better
	Sutter Memorial Hospital			3.6% (1; 35)	2.7% (1; 99)	1.5% (5; 385)	2.3% (2; 102)	6.0% (20; 432)	2.8% (11; 329)	7.7% (6; 80)	2.7% (3; 66)		<b>0.4%</b> (1; 185) Better
	UC Davis Medical Center	27.6% (1; 10)	0% (0; 17)	0% (0; 21)	5.9% (11; 260)	2.6% (18; 529)	0% (0; 42)	5.1% (23; 430)	2.8% (19; 608)	9.6% (42; 297)	1.8% (8; 299)	3.7% (4; 77)	3.5% (15; 465)
San Benito	Hazel Hawkins Memorial Hospital							0% (0; 8)	2.6% (2; 63)	9.1% (2; 34)	2.3% (1; 48)	10.4% (2; 21)	7.6% (6; 96)
San Bernardino	Arrowhead Regional Medical Center	37.1% (1; 3)	38.6% (1; 6)		<b>15.7%</b> (8; 73) Worse	8.2% (1; 76)	0% (0; 9)	9.3% (6; 121)	3.0% (8; 500)	11.7% (54; 323)	2.5% (2; 98)	0% (0; 49)	3.7% (11; 344)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

### Definition of Terms:

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	5.8% (29; 500)	5.2% (38; 732)	1.9% (50; 2,567)	6.7% (938; 13,926)	1.9% (933; 48,689)	0.7% (49; 7,411)	7.1% (3,151; 44,510)	3.3% (2,832; 85,934)	10.4% (5,285; 50,988)	2.3% (1,114; 47,720)	2.4% (571; 23,514)	4.6% (3,703; 79,958)
	Barstow Community Hospital							7.1% (2; 15)	1.6% (3; 181)	0% (0; 60) <b>Better</b>	0% (0; 33)	0% (0; 3)	0% (0; 111) <b>Better</b>
	Bear Valley Community Hospital								0% (0; 19)	0% (0; 4)	0% (0; 7)		0% (0; 41)
	Chino Valley Medical Center							5.7% (4; 22)	0.5% (2; 175) <b>Better</b>	2.4% (4; 115) <b>Better</b>	0.8% (2; 107)	0.7% (1; 38)	0.8% (2; 131) <b>Better</b>
	Colorado River Medical Center								0% (0; 14)	0% (0; 4)	0% (0; 17)		22.0% (1; 16)
	Community Hospital of San Bernardino						27.1% (1; 7) <b>Worse</b>	6.5% (5; 48)	4.4% (8; 200)	9.1% (6; 71)	0.9% (1; 90)	1.1% (1; 28)	3.4% (7; 212)
	Desert Valley Hospital							4.4% (4; 31)	0.8% (3; 224) <b>Better</b>	3.3% (7; 180) <b>Better</b>	1.2% (3; 138)	0% (0; 34)	0.8% (3; 234) <b>Better</b>
	Hi-Desert Medical Center						0% (0; 10)	8.1% (1; 10)	2.0% (2; 123)	8.6% (4; 70)	5.7% (4; 49)	4.2% (2; 47)	8.7% (13; 149) <b>Worse</b>
	Kaiser Foundation Hospital – Fontana	19.6% (1; 10)	0% (0; 7)	4.2% (1; 27)	6.2% (6; 126)		0% (0; 44)	7.0% (18; 196)	1.4% (12; 637) <b>Better</b>	9.7% (45; 502)	1.7% (9; 509)	2.7% (7; 176)	2.7% (33; 860) <b>Better</b>
	Loma Linda University Medical Center		0% (0; 9)	0% (0; 24)	9.8% (24; 309)	3.0% (7; 279)	0% (0; 21)	6.4% (20; 403)	2.8% (17; 484)	9.2% (52; 424)	2.1% (6; 187)	5.2% (4; 73)	4.9% (20; 305)
	Montclair Hospital Medical Center							5.0% (1; 8)	0.5% (1; 106) <b>Better</b>	10.9% (6; 41)	1.3% (1; 28)	0% (0; 15)	1.9% (4; 117) <b>Better</b>
	Mountains Community Hospital								23.4% (1; 6)		0% (0; 3)		0% (0; 26)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Redlands Community Hospital			17.5% (1; 13)	0% (0; 3)		0% (0; 50)	5.9% (9; 95)	<b>1.2%</b> (6; 289) Better	<b>4.7%</b> (5; 115) Better	0.5% (1; 186)	1.0% (2; 162)	3.3% (19; 444)
	Saint Bernadine Medical Center		91.2% (1; 4)	0% (0; 19)	6.1% (2; 38)	1.7% (16; 1,627)	0% (0; 39)	5.9% (38; 947)	2.5% (14; 526)	<b>5.9%</b> (12; 197) Better	1.2% (2; 175)	2.7% (2; 78)	3.1% (12; 295)
	Saint Mary Regional Medical Center			0% (0; 23)	0% (0; 7)	1.6% (4; 333)	0% (0; 45)	4.9% (22; 489)	2.0% (10; 519)	<b>5.4%</b> (8; 224) Better	1.5% (4; 218)	0.6% (1; 146)	3.0% (10; 385)
	San Antonio Community Hospital			0% (0; 6)	4.2% (3; 59)	1.7% (6; 250)	0% (0; 51)	7.9% (32; 377)	<b>1.5%</b> (11; 502) Better	<b>6.6%</b> (24; 294) Better	1.2% (4; 239)	0.9% (2; 134)	<b>2.3%</b> (15; 499) Better
	Victor Valley Community Hospital							2.8% (1; 20)	1.4% (2; 158)	7.7% (5; 63)	7.1% (3; 58)	2.3% (1; 30)	3.5% (6; 167)
San Diego	Alvarado Hospital			0% (0; 8)	0% (0; 31)	0.9% (4; 482)	0% (0; 27)	<b>2.7%</b> (4; 139) Better	2.0% (6; 257)	<b>5.7%</b> (12; 159) Better	3.8% (8; 142)	3.5% (3; 66)	5.3% (18; 358)
	Fallbrook Hospital District							0% (0; 10)	1.3% (1; 57)	0% (0; 15)	0% (0; 42)	3.8% (1; 35)	6.0% (3; 77)
	Grossmont Hospital			0% (0; 22)	5.6% (6; 114)	2.8% (22; 659)	0% (0; 102)	8.7% (43; 477)	3.2% (32; 787)	10.9% (72; 600)	2.2% (10; 365)	2.4% (8; 272)	4.5% (29; 617)
	Kaiser Foundation Hospital – San Diego	9.2% (1; 12)	0% (0; 12)	0% (0; 36)	3.4% (3; 195)		0% (0; 75)	8.2% (23; 193)	2.9% (29; 859)	8.5% (33; 484)	1.9% (8; 468)	2.7% (9; 299)	4.1% (36; 979)
	Palomar Medical Center	22.2% (2; 4)		5.5% (1; 25)	2.6% (2; 52)	2.7% (18; 409)	0% (0; 22)	8.3% (28; 361)	2.8% (20; 526)	11.7% (58; 457)	3.2% (8; 293)	4.6% (8; 162)	6.0% (31; 453)
	Paradise Valley Hospital ( <a href="#">View Comment Letter</a> )				<b>18.8%</b> (5; 7) Worse	<b>6.0%</b> (2; 6) Worse	0% (0; 7)	<b>4.1%</b> (10; 80) Better	2.4% (9; 265)	7.7% (9; 101)	0% (0; 86)	2.1% (1; 23)	<b>2.3%</b> (9; 241) Better

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Pomerado Hospital						0% (0; 8)	3.5% (4; 51)	4.8% (16; 221)	10.3% (10; 89)	0.9% (1; 123)	3.3% (3; 105)	5.0% (14; 202)
	Scripps Green Hospital		0% (0; 5)	0% (0; 41)	4.3% (2; 70)	<b>0.5%</b> (4; 1,289) <b>Better</b>	0% (0; 41)	4.0% (7; 200)	1.7% (7; 329)	<b>4.0%</b> (4; 140) <b>Better</b>	1.5% (2; 158)	1.8% (3; 108)	3.1% (11; 287)
	Scripps Memorial Hospital – Encinitas			0% (0; 13)	6.8% (2; 33)	2.8% (5; 142)	0% (0; 40)	4.5% (7; 152)	3.3% (14; 266)	7.9% (14; 200)	1.4% (3; 189)	1.1% (2; 138)	3.9% (14; 252)
	Scripps Memorial Hospital – La Jolla	0% (0; 3)	13.6% (1; 6)	<b>22.4%</b> (2; 18) <b>Worse</b>	8.5% (7; 129)	2.0% (15; 1,266)	0% (0; 19)	8.3% (28; 478)	1.9% (7; 270)	8.7% (21; 248)	1.1% (2; 164)	1.3% (3; 133)	4.3% (10; 224)
	Scripps Mercy Hospital		0% (0; 4)	0% (0; 13)	7.4% (8; 57)	2.0% (11; 487)	0% (0; 42)	7.1% (32; 406)	<b>1.7%</b> (16; 776) <b>Better</b>	<b>8.6%</b> (65; 509) <b>Better</b>	1.8% (9; 406)	1.3% (3; 184)	<b>2.0%</b> (14; 590) <b>Better</b>
	Sharp Chula Vista Medical Center	0% (0; 3)		0% (0; 8)	11.6% (8; 44)	1.0% (6; 368)	0% (0; 21)	<b>4.2%</b> (16; 305) <b>Better</b>	2.8% (20; 541)	10.2% (34; 285)	1.2% (4; 248)	4.2% (3; 79)	3.6% (16; 424)
	Sharp Coronado Hospital and Healthcare Center							0% (0; 6)	2.9% (4; 70)	7.7% (2; 22)	0% (0; 41)	3.8% (2; 33)	4.1% (3; 68)
	Sharp Memorial Hospital ( <a href="#">View Comment Letter</a> )	14.3% (1; 7)	0% (0; 5)	0% (0; 25)	<b>12.0%</b> (12; 66) <b>Worse</b>	1.5% (6; 333)	1.5% (1; 56)	6.7% (22; 378)	<b>5.3%</b> (29; 513) <b>Worse</b>	11.9% (49; 387)	1.4% (6; 362)	1.2% (3; 240)	<b>7.8%</b> (37; 524) <b>Worse</b>
	Tri-City Medical Center – Oceanside			0% (0; 19)	<b>15.0%</b> (7; 43) <b>Worse</b>	2.3% (5; 291)	0% (0; 38)	9.5% (36; 323)	3.0% (14; 448)	11.4% (50; 383)	1.8% (5; 281)	3.0% (5; 172)	4.4% (27; 545)
	UC San Diego Medical Center	0% (0; 7)	14.8% (3; 31)	13.9% (1; 10)	6.3% (20; 235)	<b>0.3%</b> (1; 366) <b>Better</b>	7.6% (1; 29)	<b>3.6%</b> (9; 257) <b>Better</b>	3.4% (14; 382)	8.6% (38; 313)	<b>0.3%</b> (1; 228) <b>Better</b>	0% (0; 94)	<b>2.6%</b> (11; 406) <b>Better</b>

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
San Francisco	California Pacific Medical Center – Pacific Campus	0% (0; 10)	0% (0; 38)	8.4% (2; 24)	5.0% (14; 198)	2.2% (12; 348)	2.3% (1; 44)	8.2% (24; 245)	3.5% (25; 552)	<b>8.0%</b> (56; 467) <b>Better</b>	3.2% (15; 362)	1.5% (3; 142)	<b>2.6%</b> (11; 381) <b>Better</b>
	California Pacific Medical Center – St. Luke’s Campus						0% (0; 6)	6.9% (2; 16)	1.7% (4; 213)	10.3% (3; 28)	5.7% (2; 71)	6.2% (2; 25)	2.1% (3; 133)
	Chinese Hospital		14.7% (1; 3)					0% (0; 20)	0.9% (1; 90)	8.5% (2; 57)	3.2% (2; 76)	0% (0; 44)	7.8% (10; 174)
	Kaiser Foundation Hospital – San Francisco		0% (0; 17)	3.3% (1; 57)		2.0% (15; 872)	1.2% (1; 70)	<b>4.1%</b> (16; 682) <b>Better</b>	<b>1.2%</b> (6; 382) <b>Better</b>	7.5% (17; 201)	2.6% (5; 188)	4.4% (3; 69)	3.8% (10; 245)
	Laguna Honda Hospital and Rehabilitation Center												11.9% (3; 28)
	Saint Francis Memorial Hospital				4.4% (1; 15)			<b>14.0%</b> (9; 31) <b>Worse</b>	<b>0.8%</b> (2; 202) <b>Better</b>	7.7% (17; 137)	3.7% (5; 77)	2.6% (1; 56)	6.7% (15; 196)
	Saint Mary’s Medical Center, San Francisco			0% (0; 4)	0% (0; 3)	2.2% (5; 287)	0% (0; 13)	7.3% (10; 127)	2.5% (11; 252)	6.2% (7; 95)	2.3% (4; 125)	4.8% (4; 47)	3.5% (14; 217)
	San Francisco General Hospital				10.7% (8; 56)	<b>10.3%</b> (4; 87) <b>Worse</b>		10.2% (8; 138)	2.8% (5; 313)	13.4% (33; 232)	3.0% (6; 164)	4.8% (2; 25)	2.0% (4; 391)
	UC San Francisco Medical Center	0% (0; 26)	7.3% (1; 24)	0% (0; 49)	5.4% (55; 1,245)	3.2% (7; 195)	0% (0; 53)	8.3% (14; 161)	3.9% (23; 453)	10.2% (95; 521)	2.0% (6; 200)	1.6% (2; 61)	3.6% (18; 453)
San Joaquin	Dameron Hospital			0% (0; 3)		3.3% (2; 172)	0% (0; 13)	5.9% (11; 277)	4.1% (10; 273)	8.2% (7; 122)	2.8% (4; 176)	<b>6.8%</b> (6; 76) <b>Worse</b>	<b>8.4%</b> (29; 312) <b>Worse</b>
	Doctors Hospital of Manteca							20.2% (2; 9)	1.8% (3; 150)	15.0% (10; 55)	1.7% (1; 73)	0% (0; 27)	1.4% (1; 97)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Kaiser Foundation Hospital – Manteca			0% (0; 17)			0% (0; 53)	4.1% (3; 47)	3.5% (9; 247)	12.2% (10; 135)	0% (0; 153)	1.5% (2; 83)	5.1% (16; 290)
	Lodi Memorial Hospital	0% (0; 3)	0% (0; 3)	20.8% (1; 8)			0% (0; 28)	7.6% (7; 58)	1.5% (3; 206)	10.7% (15; 129)	2.5% (3; 140)	1.1% (1; 86)	<b>1.5%</b> <b>(4; 323)</b> <b>Better</b>
	Saint Joseph’s Medical Center of Stockton	0% (0; 4)		0% (0; 23)	6.2% (3; 56)	1.5% (5; 511)	1.7% (1; 123)	5.7% (18; 348)	3.3% (23; 549)	9.3% (39; 290)	2.8% (9; 313)	1.5% (2; 137)	5.1% (34; 560)
	San Joaquin General Hospital				7.1% (2; 12)		0% (0; 7)	11.7% (5; 63)	1.9% (4; 238)	10.1% (21; 122)	5.5% (5; 130)	5.0% (2; 27)	3.8% (6; 175)
	Sutter Tracy Community Hospital							14.7% (4; 17)	1.4% (2; 117)	7.4% (4; 53)	2.2% (1; 63)	0% (0; 48)	3.7% (4; 142)
San Luis Obispo	Arroyo Grande Community Hospital						0% (0; 20)	12.2% (6; 28)	5.3% (6; 101)	9.4% (10; 67)	3.4% (2; 73)	4.4% (3; 48)	3.8% (3; 82)
	French Hospital Medical Center			0% (0; 3)		2.7% (7; 380)	0% (0; 38)	6.8% (9; 205)	2.9% (6; 128)	17.8% (7; 78)	2.6% (3; 78)	3.0% (1; 42)	3.7% (6; 158)
	Sierra Vista Regional Medical Center ( <a href="#">View Comment Letter</a> )			0% (0; 19)	11.1% (7; 106)	5.4% (2; 15)	0% (0; 16)	8.6% (4; 31)	3.0% (2; 77)	<b>15.9%</b> <b>(31; 173)</b> <b>Worse</b>	0% (0; 79)	0% (0; 51)	5.9% (6; 163)
	Twin Cities Community Hospital						0% (0; 5)	15.7% (6; 22)	3.3% (5; 162)	14.7% (14; 127)	1.0% (1; 126)	1.4% (1; 90)	6.5% (13; 242)
San Mateo	Kaiser Foundation Hospital – Redwood City ( <a href="#">View Comment Letter</a> )				6.0% (22; 652)			7.1% (7; 47)	2.2% (6; 146)	<b>13.8%</b> <b>(65; 395)</b> <b>Worse</b>	2.0% (2; 108)	1.2% (1; 91)	<b>7.8%</b> <b>(21; 214)</b> <b>Worse</b>
	Kaiser Foundation Hospital – South San Francisco						0% (0; 6)	6.2% (6; 47)	3.5% (10; 173)	9.2% (14; 136)	1.8% (4; 180)	1.8% (2; 72)	4.1% (17; 276)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurysm Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Peninsula Medical Center	0% (0; 3)	31.7% (2; 9)	0% (0; 19)	10.5% (4; 16)	1.6% (5; 244)	0% (0; 70)	9.0% (17; 180)	2.2% (13; 321)	8.6% (28; 259)	1.0% (3; 216)	1.2% (2; 154)	3.6% (15; 317)
	San Mateo Medical Center				0% (0; 6)			13.2% (5; 27)	2.6% (3; 117)	11.1% (3; 41)	0% (0; 67)	0% (0; 17)	<b>11.1%</b> (11; 140) Worse
	Sequoia Hospital			0% (0; 13)		1.7% (2; 164)	0% (0; 40)	11.6% (7; 88)	3.0% (10; 260)	13.6% (11; 77)	1.1% (1; 96)	1.7% (2; 86)	3.5% (7; 167)
	Seton Medical Center			0% (0; 3)	8.4% (2; 14)	1.2% (4; 428)	0% (0; 27)	7.2% (9; 141)	2.5% (17; 541)	9.4% (22; 181)	<b>5.0%</b> (13; 205) Worse	2.9% (2; 83)	4.1% (17; 283)
Santa Barbara	Goleta Valley Cottage Hospital								0% (0; 33)	22.2% (1; 4)	0% (0; 12)	0% (0; 58)	0% (0; 47)
	Lompoc Valley Medical Center						0% (0; 3)	9.6% (2; 10)	4.2% (4; 85)	9.9% (5; 64)	0% (0; 61)	7.8% (2; 31)	<b>12.7%</b> (13; 105) Worse
	Marian Medical Center			0% (0; 9)	11.9% (2; 29)	0.9% (2; 143)	0% (0; 66)	7.9% (18; 155)	2.1% (11; 269)	11.1% (20; 175)	1.9% (4; 189)	0.9% (1; 96)	3.2% (11; 185)
	Santa Barbara Cottage Hospital	0% (0; 5)	0% (0; 7)	9.9% (1; 23)	5.7% (17; 246)	3.1% (10; 455)	4.0% (1; 56)	8.3% (16; 231)	3.5% (14; 347)	10.2% (52; 467)	3.8% (10; 276)	5.4% (7; 142)	5.0% (16; 346)
	Santa Ynez Valley Cottage Hospital							0% (0; 5)	6.2% (1; 18)	0% (0; 3)	0% (0; 5)		5.1% (1; 33)
Santa Clara	El Camino Hospital			6.4% (2; 57)	7.1% (5; 43)	2.7% (12; 228)	0% (0; 36)	<b>11.7%</b> (29; 209) Worse	3.8% (20; 372)	<b>15.6%</b> (47; 263) Worse	4.4% (9; 243)	1.4% (2; 171)	5.0% (20; 380)
	El Camino Hospital Los Gatos**							0% (0; 4)	0% (0; 20)	<b>0%</b> (0; 7) Better	0% (0; 12)	0% (0; 6)	3.5% (1; 24)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Good Samaritan Hospital – San Jose		0% (0; 3)	0% (0; 13)	9.1% (15; 103)	1.9% (10; 269)	0% (0; 34)	5.9% (16; 248)	2.6% (17; 370)	9.1% (40; 326)	2.3% (8; 207)	3.9% (5; 144)	5.6% (22; 314)
	Kaiser Foundation Hospital – San Jose		0% (0; 6)	0% (0; 5)		3.1% (4; 62)	0% (0; 33)	8.6% (16; 158)	2.6% (12; 325)	12.5% (34; 275)	1.9% (4; 204)	5.3% (6; 108)	<b>6.9%</b> (29; 374) Worse
	Kaiser Foundation Hospital – Santa Clara	0% (0; 7)		3.2% (1; 36)		1.8% (13; 1,270)	0% (0; 47)	4.8% (22; 790)	3.4% (28; 594)	13.0% (43; 290)	1.7% (5; 279)	4.2% (7; 143)	6.3% (29; 398)
	O’Connor Hospital – San Jose			4.4% (1; 21)	7.6% (3; 44)	2.2% (5; 280)	0% (0; 33)	9.5% (13; 159)	2.3% (12; 434)	9.3% (19; 182)	2.6% (4; 206)	1.0% (1; 97)	<b>2.7%</b> (13; 364) Better
	Regional Medical of San Jose				5.9% (8; 44)	2.0% (9; 265)	<b>16.0%</b> (2; 13) Worse	8.1% (19; 193)	3.1% (16; 396)	8.3% (37; 295)	0.8% (2; 207)	3.1% (4; 76)	3.4% (12; 326)
	Saint Louise Regional Hospital							11.8% (3; 22)	1.7% (2; 106)	9.4% (2; 24)	1.1% (1; 74)	0% (0; 35)	<b>9.3%</b> (12; 136) Worse
	Santa Clara Valley Medical Center		0% (0; 3)		10.6% (7; 68)	2.2% (5; 206)	0% (0; 6)	9.0% (12; 198)	3.1% (11; 438)	8.2% (19; 276)	2.2% (6; 271)	0% (0; 42)	3.3% (11; 428)
	Stanford Hospital	0% (0; 35)	6.6% (2; 39)	0% (0; 64)	<b>3.9%</b> (34; 830) Better	2.2% (8; 328)	0% (0; 82)	6.0% (15; 207)	4.3% (34; 453)	<b>7.7%</b> (58; 500) Better	1.5% (5; 254)	2.2% (4; 121)	4.1% (20; 432)
Santa Cruz	Dominican Hospital – Santa Cruz/Soquel			8.6% (1; 6)	11.0% (7; 61)	2.4% (7; 288)	0% (0; 23)	8.3% (20; 223)	3.1% (14; 304)	12.5% (24; 227)	2.7% (6; 203)	1.9% (3; 124)	4.6% (13; 272)
	Watsonville Community Hospital		15.4% (1; 3)				0% (0; 3)	13.9% (7; 28)	4.9% (12; 209)	9.7% (6; 70)	1.3% (1; 110)	5.1% (2; 35)	2.0% (2; 133)
Shasta	Mayers Memorial Hospital								8.8% (1; 16)	63.0% (1; 3)	0% (0; 5)	0% (0; 5)	0% (0; 28)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurysm Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Mercy Medical Center – Redding			0% (0; 34)	5.9% (5; 66)	2.5% (11; 405)	1.2% (1; 147)	7.6% (27; 362)	3.6% (22; 369)	<b>7.3%</b> (23; 263) <b>Better</b>	2.0% (6; 203)	<b>0.3%</b> (1; 174) <b>Better</b>	<b>2.0%</b> (11; 407) <b>Better</b>
	Shasta Regional Medical Center ( <a href="#">View Comment Letter</a> )					0.6% (1; 58)	5.6% (1; 16)	5.4% (6; 72)	<b>1.3%</b> (4; 139) <b>Better</b>	5.3% (4; 105)	0.8% (1; 103)	2.2% (2; 34)	<b>0.9%</b> (3; 180) <b>Better</b>
Siskiyou	Fairchild Medical Center						0% (0; 8)		7.5% (5; 45)	0% (0; 21)	4.6% (2; 56)	0% (0; 15)	4.3% (5; 145)
	Mercy Medical Center Mt. Shasta						0% (0; 3)	5.3% (1; 11)	6.3% (2; 39)	11.9% (4; 14)	7.5% (1; 18)	0% (0; 15)	7.3% (4; 74)
Solano	Kaiser Foundation Hospital – Rehabilitation Center Vallejo			0% (0; 10)			0% (0; 64)	4.6% (8; 102)	2.3% (13; 406)	<b>6.5%</b> (19; 316) <b>Better</b>	1.3% (4; 260)	2.3% (4; 141)	6.0% (42; 573)
	Kaiser Foundation Hospital – Vacaville							5.7% (1; 13)	6.9% (2; 32)	16.5% (4; 21)	8.9% (1; 14)	0% (0; 8)	2.3% (1; 52)
	North Bay Medical Center					0% (0; 21)		7.2% (6; 57)	2.6% (4; 168)	7.9% (4; 55)	0% (0; 85)	0% (0; 35)	4.7% (11; 215)
	North Bay VacaValley Hospital							9.7% (6; 23)	<b>0%</b> (0; 111) <b>Better</b>	12.8% (8; 55)	2.0% (2; 77)	2.3% (1; 37)	4.4% (11; 170)
	Sutter Solano Medical Center						0% (0; 6)	5.0% (2; 23)	<b>0.6%</b> (1; 173) <b>Better</b>	<b>3.9%</b> (4; 84) <b>Better</b>	4.5% (6; 88)	7.1% (2; 32)	3.8% (4; 103)
Sonoma	Healdsburg District Hospital							0% (0; 8)	3.4% (1; 23)	7.3% (1; 25)	0% (0; 5)	4.2% (1; 22)	1.3% (1; 58)
	Kaiser Foundation Hospital – Santa Rosa						0% (0; 33)	4.5% (5; 83)	3.8% (7; 156)	10.7% (21; 186)	0.6% (1; 166)	1.6% (3; 120)	5.6% (17; 301)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

### Definition of Terms:

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	<b>5.8%</b> (29; 500)	<b>5.2%</b> (38; 732)	<b>1.9%</b> (50; 2,567)	<b>6.7%</b> (938; 13,926)	<b>1.9%</b> (933; 48,689)	<b>0.7%</b> (49; 7,411)	<b>7.1%</b> (3,151; 44,510)	<b>3.3%</b> (2,832; 85,934)	<b>10.4%</b> (5,285; 50,988)	<b>2.3%</b> (1,114; 47,720)	<b>2.4%</b> (571; 23,514)	<b>4.6%</b> (3,703; 79,958)
	Palm Drive Hospital							0% (0; 8)	0% (0; 33)	3.5% (1; 20)	3.1% (1; 33)	0% (0; 28)	4.8% (6; 94)
	Petaluma Valley Hospital							6.9% (9; 55)	0.8% (1; 85)	12.4% (6; 50)	3.9% (2; 51)	4.2% (3; 45)	3.1% (5; 107)
	Santa Rosa Memorial Hospital – Montgomery			0% (0; 14)	14.9% (5; 20)	2.9% (9; 211)	0% (0; 42)	7.8% (20; 227)	3.6% (22; 313)	12.8% (33; 226)	3.4% (9; 198)	2.2% (6; 167)	5.0% (16; 231)
	Sonoma Valley Hospital						0% (0; 3)	3.1% (1; 14)	0% (0; 53)	6.5% (4; 59)	0% (0; 36)	0% (0; 38)	2.8% (2; 57)
	Sutter Medical Center of Santa Rosa			0% (0; 12)		1.5% (3; 233)	0% (0; 56)	6.4% (6; 159)	1.5% (2; 115)	12.1% (3; 38)	1.2% (1; 68)	5.4% (2; 26)	4.0% (3; 74)
Stanislaus	Doctors Medical Center		0% (0; 3)	9.3% (2; 26)	5.9% (3; 39)	1.2% (6; 554)	0% (0; 136)	7.0% (29; 460)	2.5% (12; 457)	9.1% (28; 224)	1.5% (4; 171)	3.2% (5; 104)	4.5% (17; 380)
	Emanuel Medical Center, Inc.						0% (0; 25)	5.4% (9; 75)	3.0% (13; 341)	11.7% (15; 142)	2.6% (5; 184)	0% (0; 71)	3.2% (16; 443)
	Memorial Hospital Medical Center – Modesto		0% (0; 3)	9.0% (1; 13)	7.6% (4; 27)	1.5% (11; 776)	1.3% (2; 149)	<b>4.2%</b> (21; 458) <b>Better</b>	2.6% (24; 607)	<b>7.7%</b> (32; 337) <b>Better</b>	1.9% (9; 335)	1.1% (4; 252)	<b>3.1%</b> (26; 686) <b>Better</b>
	Oak Valley District Hospital							14.4% (2; 9)	4.3% (3; 42)	0% (0; 20)	0% (0; 30)	1.8% (1; 24)	4.0% (4; 65)
Sutter	Fremont Medical Center								7.4% (7; 77)	14.9% (5; 46)	3.6% (2; 65)	0% (0; 22)	<b>9.5%</b> (15; 216) <b>Worse</b>
Tehama	Saint Elizabeth Community Hospital							7.0% (4; 15)	1.8% (2; 99)	8.6% (4; 76)	0% (0; 51)	4.5% (1; 24)	2.8% (4; 148)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	<b>STATEWIDE</b>	5.8% (29; 500)	5.2% (38; 732)	1.9% (50; 2,567)	6.7% (938; 13,926)	1.9% (933; 48,689)	0.7% (49; 7,411)	7.1% (3,151; 44,510)	3.3% (2,832; 85,934)	10.4% (5,285; 50,988)	2.3% (1,114; 47,720)	2.4% (571; 23,514)	4.6% (3,703; 79,958)
Trinity	Trinity Hospital							41.3% (1; 4)	0% (0; 17)	0% (0; 4)	0% (0; 11)		2.6% (1; 51)
Tulare	Kaweah Delta Medical Center			0% (0; 16)	14.9% (1; 4)	1.8% (14; 1,194)	3.3% (1; 105)	6.9% (21; 386)	3.1% (11; 361)	<b>16.2%</b> <b>(36; 275)</b> Worse	<b>4.8%</b> <b>(12; 253)</b> Worse	4.1% (6; 161)	4.8% (26; 587)
	Sierra View District Hospital							<b>17.1%</b> <b>(10; 61)</b> Worse	<b>6.1%</b> <b>(18; 259)</b> Worse	8.3% (9; 127)	2.3% (4; 136)	0% (0; 61)	<b>9.0%</b> <b>(36; 395)</b> Worse
	Tulare District Hospital							9.2% (3; 22)	4.6% (4; 123)	12.3% (6; 74)	2.4% (1; 64)	3.6% (2; 29)	3.6% (5; 233)
Tuolumne	Sonora Regional Medical Center – Greenley						0% (0; 14)	5.4% (5; 46)	2.8% (3; 77)	16.6% (8; 64)	1.4% (2; 92)	2.3% (2; 84)	2.9% (6; 191)
Ventura	Community Memorial Hospital – San Buenaventura			0% (0; 14)	8.4% (2; 31)	2.5% (10; 365)	4.6% (2; 68)	6.8% (15; 266)	2.4% (11; 333)	10.3% (24; 218)	<b>0.7%</b> <b>(2; 210)</b> Better	1.4% (2; 128)	4.8% (20; 283)
	Los Robles Hospital and Medical Center			0% (0; 22)	13.1% (3; 39)	1.3% (4; 408)	0% (0; 25)	6.1% (16; 277)	2.0% (8; 293)	9.8% (20; 207)	3.1% (6; 179)	1.9% (3; 156)	3.2% (8; 230)
	Ojai Valley Community Hospital							13.6% (3; 14)	11.9% (2; 16)	5.9% (1; 17)	0% (0; 8)	0% (0; 24)	0% (0; 35)
	Saint John’s Pleasant Valley Hospital							5.3% (3; 27)	2.1% (6; 129)	12.5% (13; 88)	2.2% (2; 85)	0% (0; 67)	2.8% (5; 165)
	Saint John’s Regional Medical Center			3.9% (1; 17)	5.4% (2; 20)	1.8% (10; 263)	1.7% (1; 52)	<b>3.6%</b> <b>(12; 257)</b> Better	<b>1.3%</b> <b>(7; 365)</b> Better	8.9% (28; 216)	0.8% (2; 177)	0% (0; 81)	3.6% (18; 320)
	Simi Valley Hospital and Healthcare Services – Sycamore				0% (0; 5)		0% (0; 7)	3.4% (3; 38)	3.4% (5; 125)	7.7% (10; 104)	1.1% (1; 87)	0% (0; 60)	4.5% (9; 200)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.

## California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2009

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Transluminal Coronary Angioplasty	Carotid Endarterectomy	Acute Myocardial Infarction	Congestive Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths and total cases, and the third line indicates any quality rating.													
	STATEWIDE	5.8% (29; 500)	5.2% (38; 732)	1.9% (50; 2,567)	6.7% (938; 13,926)	1.9% (933; 48,689)	0.7% (49; 7,411)	7.1% (3,151; 44,510)	3.3% (2,832; 85,934)	10.4% (5,285; 50,988)	2.3% (1,114; 47,720)	2.4% (571; 23,514)	4.6% (3,703; 79,958)
	Thousand Oaks Surgical Hospital				0% (0; 40)								
	Ventura County Medical Center	0% (0; 3)			0% (0; 31)			11.7% (4; 35)	3.5% (5; 149)	9.3% (9; 122)	2.1% (2; 99)	0% (0; 39)	7.2% (12; 184)
Yolo	Sutter Davis Hospital						0% (0; 6)	13.4% (2; 4)	0.9% (1; 58)	5.7% (1; 29)	0% (0; 42)	0% (0; 46)	4.9% (6; 104)
	Woodland Memorial Hospital						0% (0; 10)	9.2% (5; 30)	1.3% (1; 60)	8.4% (9; 78)	3.8% (5; 64)	2.9% (1; 47)	3.0% (6; 115)
Yuba	Rideout Memorial Hospital	11.4% (1; 3)		0% (0; 10)	7.7% (2; 17)	4.2% (10; 353) Worse	0% (0; 68)	5.5% (11; 216)	4.9% (16; 320)	9.4% (13; 144)	3.7% (5; 140)	4.9% (5; 111)	4.4% (9; 295)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.1, and OSHPD hospital patient discharge data, 2009. See [Technical Note](#) for details about methodology.

\*\*Hospital name change from 2008 to 2009, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2008 and 2009 Data, [Technical Note](#).

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

**Definition of Terms:**

**Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to “level the playing field” and allow fair comparisons across hospitals.

**Number of Deaths:** number of patients that died in this hospital.

**Number of Cases:** total number of patients undergoing this procedure or with this medical condition in this hospital.

**Quality Ratings** are risk-adjusted using age, gender APR-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate (“better”) or lower 95% confidence level of the RAMR is higher than the statewide rate (“worse”). Cells with no quality ratings indicate a hospital’s performance is considered “not significantly different” from the statewide average.