

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
Alameda	Alameda County Medical Center – Highland Campus				9.8% (2;32)		0.0% (0;3)	12.7% (3;29)	2.7% (3;339)	11.8% (12;152)	3.4% (2;141)	0.0% (0;13)	5.7% (4;212)
	Alameda Hospital						0.0% (0;3)	5.3% (6;56)	2.5% (5;162)	17.1% (9;53)	5.3% (3;70)	2.2% (1;43)	1.9% (4;150)
	Alta Bates Summit Medical Center – Alta Bates Campus				5.7% (4;48)	2.6% (5;95)	6.9% (1;13)	6.0% (8;106)	2.6% (8;286)	7.8% (23;251)	0.9% (2;196)	1.5% (2;104)	3.3% (8;245)
	Alta Bates Summit Medical Center – Summit Campus – Hawthorne	0.0% (0;3)	0.0% (0;3)	0.0% (0;6)	41.7% (1;5)	2.9% (19;792)	7.2% (1;21)	6.9% (24;525)	2.0% (10;524)	9.1% (21;290)	2.2% (7;297)	3.5% (4;103)	2.4% (10;371)
	Eden Medical Center				2.4% (6;117) Better		0.0% (0;7)	4.0% (5;49)	2.8% (9;223)	8.3% (63;384) Better	1.6% (3;159)	0.0% (0;66)	3.4% (8;195)
	Kaiser Foundation Hospital – Hayward		12.2% (1;10)	0.0% (0;16)		0.0% (0;6)	0.0% (0;38)	4.6% (10;153)	3.4% (16;454)	8.0% (18;276)	3.1% (10;332)	4.5% (6;130)	4.0% (17;378)
	Kaiser Foundation Hospital – Oakland Campus		0.0% (0;10)	0.0% (0;16)	0.0% (0;5)		0.0% (0;32)	3.8% (6;106)	2.2% (11;590)	9.3% (22;348)	4.2% (8;271)	5.4% (5;124)	4.0% (14;377)
	Saint Rose Hospital					3.4% (7;113)	0.0% (0;3)	12.5% (16;144) Worse	2.2% (4;273)	2.9% (1;76)	4.4% (5;154)	0.0% (0;30)	0.0% (0;132)
	San Leandro Hospital			0.0% (0;20)			0.0% (0;23)	8.6% (8;34)	2.5% (6;271)	6.1% (2;55)	0.9% (1;122)	2.3% (1;41)	4.0% (4;110)
Valleycare Medical Center			0.0% (0;5)		2.8% (4;125)	0.0% (0;31)	11.5% (16;150) Worse	3.5% (10;216)	14.5% (12;88)	2.1% (3;107)	4.2% (2;50)	4.8% (14;234)	
Washington Hospital – Fremont				100% (1;5) Worse	8.3% (8;57)	2.3% (11;357)	0.0% (0;16)	9.2% (21;281)	4.5% (16;381)	13.4% (37;279) Worse	3.2% (9;249)	0.0% (0;100)	6.2% (25;375) Worse
Amador	Sutter Amador Hospital							6.6% (1;10)	4.8% (4;80)	7.8% (3;54)	2.5% (1;43)	2.5% (1;52)	2.2% (2;109)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia		
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.															
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)		
Butte	Biggs Gridley Memorial Hospital								10.1% (1;19)	55.9% (2;6)	0.0% (0;17)	0.0% (0;14)	9.2% (3;88)		
	Enloe Medical Center – Esplanade Campus			0.0% (0;19)	13.3% (6;43)	2.0% (7;394)	0.0% (0;85)	8.1% (23;309)	3.5% (11;239)	14.6% (31;228) Worse	2.3% (8;275)	3.9% (7;177)	5.0% (19;460)		
	Feather River Hospital						0.0% (0;18)	1.2% (1;42) Better	3.3% (6;114)	6.6% (4;77)	1.2% (2;101)	0.8% (1;63)	1.9% (4;166)		
	Oroville Hospital						0.0% (0;11)	4.9% (5;48)	1.1% (4;294) Better	6.5% (7;117)	1.1% (2;164)	1.7% (2;64)	1.8% (7;417) Better		
Calaveras	Mark Twain Saint Joseph’s Hospital							25.6% (3;9)	0.0% (0;46)	20.9% (7;42) Worse	3.7% (1;32)	0.0% (0;26)	2.2% (1;79)		
Colusa	Colusa Regional Medical Center							0.0% (0;5)	4.4% (1;28)	33.9% (2;8)	6.4% (1;26)		6.5% (3;60)		
Contra Costa	Contra Costa Regional Medical Center							8.8% (2;20)	1.3% (2;282)	11.0% (2;69)	0.0% (0;70)	0.0% (0;23)	4.6% (6;234)		
	Doctors Medical Center – San Pablo							5.5% (6;158) Worse	0.0% (0;4)	6.8% (12;182)	3.3% (12;462)	8.6% (11;131)	4.1% (5;132)	4.5% (2;49)	6.6% (13;200)
	John Muir Medical Center – Concord Campus	8.8% (1;9)		0.0% (0;44)		0.9% (3;238)	0.0% (0;45)	7.1% (19;285)	2.4% (12;386)	18.1% (23;138) Worse	3.0% (7;178)	1.4% (1;83)	6.2% (16;225)		
	John Muir Medical Center – Walnut Creek Campus		0.0% (0;3)		2.7% (6;114) Better	1.6% (4;172)	0.0% (0;45)	5.2% (17;205)	3.4% (17;327)	10.0% (64;374)	0.3% (1;249) Better	2.7% (7;198)	3.2% (12;293)		
	Kaiser Foundation Hospital – Antioch							5.8% (8;85)	3.6% (10;253)	8.2% (12;141)	1.8% (2;129)	3.0% (2;66)	3.5% (7;230)		

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Kaiser Foundation Hospital – Walnut Creek	0.0% (0;9)	0.0% (0;13)	2.5% (1;38)		1.7% (3;154)	2.8% (1;48)	6.5% (9;166)	2.9% (13;427)	12.8% (25;251)	1.3% (3;251)	4.5% (7;210)	5.9% (17;344)
	San Ramon Regional Medical Center			0.0% (0;3)		3.6% (2;50)	0.0% (0;10)	9.7% (5;53)	3.5% (3;91)	14.9% (7;58)	5.5% (3;74)	8.2% (3;44)	6.1% (7;139)
	Sutter Delta Medical Center					0.6% (1;84)	0.0% (0;10)	4.3% (8;152)	1.3% (5;294) Better	11.9% (9;97)	0.6% (1;148)	0.0% (0;56)	2.2% (7;259)
Del Norte	Sutter Coast Hospital							0.0% (0;20)	7.0% (7;75)	17.7% (8;52)	3.2% (2;48)		2.6% (4;117)
El Dorado	Barton Memorial Hospital							55.4% (1;3)	2.6% (1;38)	42.8% (3;19)	1.9% (1;39)	9.0% (1;21)	4.9% (2;86)
	Marshall Medical Center						0.0% (0;7)	9.9% (4;27)	3.6% (7;142)	6.5% (5;85)	2.1% (3;104)	0.8% (1;59)	1.5% (4;189) Better
Fresno	Clovis Community Medical Center		0.0% (0;6)			0.0% (0;57)		6.5% (4;88)	4.5% (5;131)	10.4% (3;87)	2.3% (2;116)	3.4% (2;82)	5.2% (13;292)
	Coalinga Regional Medical Center							0.0% (0;4)	5.7% (1;28)	0.0% (0;5)	0.0% (0;6)		0.0% (0;155)
	Community Regional Medical Center – Fresno		18.0% (1;8)	0.0% (0;11)	7.6% (22;257)	4.7% (27;580) Worse	1.3% (1;30)	9.2% (50;700) Worse	3.6% (20;803)	10.9% (123;798)	2.3% (12;533)	3.0% (5;141)	4.3% (19;607)
	Fresno Heart and Surgical Hospital	0.0% (0;3)	25.3% (2;7)	0.0% (0;18)		3.8% (4;505)	0.0% (0;125)	4.5% (2;80)	8.8% (9;140) Worse	0.0% (0;4)	0.0% (0;16)		0.0% (0;11)
	Kaiser Foundation Hospital – Fresno			0.0% (0;6)	0.0% (0;3)	0.0% (0;3)	0.0% (0;22)	5.9% (9;108)	2.6% (7;265)	10.5% (18;193)	1.0% (1;115)	4.5% (3;107)	4.0% (12;318)
	Saint Agnes Medical Center		0.0% (0;3)	0.0% (0;19)	0.0% (0;13)	2.3% (13;641)	1.7% (1;53)	6.3% (32;530)	2.9% (29;959)	11.6% (36;455)	2.2% (16;577)	4.8% (11;284) Worse	6.6% (45;792) Worse

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Sierra Kings District Hospital							23.1% (1;4)	0.0% (0;34)	30.3% (2;7)	24.3% (1;9)		15.6% (5;67) Worse
Glenn	Glenn Medical Center								0.0% (0;10)	0.0% (0;4)	0.0% (0;7)		0.0% (0;39)
Humboldt	Jerold Phelps Community Hospital												0.0% (0;8)
	Mad River Community Hospital						0.0% (0;3)	12.9% (2;7)	5.9% (2;30)	43.6% (4;36) Worse	8.0% (2;41)	7.7% (1;13)	5.3% (5;118)
	Redwood Memorial Hospital							16.2% (2;8)	6.7% (3;41)	6.2% (1;22)	7.0% (1;26)	0.0% (0;11)	1.7% (1;82)
	Saint Joseph Hospital – Eureka			0.0% (0;16)	5.4% (2;21)	3.4% (5;168)	0.0% (0;102)	12.2% (11;135)	3.2% (4;109)	12.6% (13;111)	1.7% (2;100)	2.2% (2;91)	2.5% (4;136)
Imperial	El Centro Regional Medical Center				0.0% (0;8)			6.8% (4;50)	2.6% (5;207)	14.0% (13;126)	0.0% (0;151)	6.6% (4;75)	6.1% (14;203)
	Pioneers Memorial Hospital				6.9% (1;9)			1.9% (1;35)	1.6% (2;157)	0.0% (0;83) Better	2.3% (3;133)	0.0% (0;28)	5.0% (10;220)
Inyo	Northern Inyo Hospital							40.0% (2;4)	10.1% (2;25)	31.3% (1;7)	0.0% (0;18)	5.7% (1;18)	4.0% (1;39)
	Southern Inyo Hospital												0.0% (0;4)
Kern	Bakersfield Heart Hospital			4.6% (1;10)		0.8% (4;500) Better	0.0% (0;7)	5.2% (17;392)	1.7% (7;336)	4.7% (1;45)	1.4% (2;79)		1.2% (2;151) Better
	Bakersfield Memorial Hospital			0.0% (0;31)	7.8% (7;68)	2.3% (10;376)	0.0% (0;82)	6.1% (21;294)	2.1% (8;420)	7.6% (20;242)	2.7% (8;231)	1.5% (2;119)	4.4% (18;407)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Delano Regional Medical Center							18.6% (3;10)	5.8% (6;127)	15.7% (2;20)	13.0% (3;40) Worse	14.2% (1;18)	9.9% (16;140) Worse
	Good Samaritan Hospital – Bakersfield								0.0% (0;37)		0.0% (0;11)		0.0% (0;81)
	Kern Medical Center				10.8% (3;29)		0.0% (0;4)	4.5% (1;21)	3.7% (2;103)	9.8% (5;64)	0.0% (0;85)	0.0% (0;10)	10.4% (6;117)
	Kern Valley Healthcare District							0.0% (0;6)	0.0% (0;48)	0.0% (0;10)	0.0% (0;13)		1.6% (1;124)
	Mercy Hospital – Bakersfield			0.0% (0;17)	2.1% (1;28)		0.0% (0;34)	12.5% (9;29)	6.4% (11;171) Worse	7.3% (9;134)	2.6% (6;202)	0.0% (0;100)	4.6% (18;493)
	Ridgecrest Regional Hospital							22.9% (2;5)	4.9% (3;76)	16.0% (2;23)	11.9% (4;59) Worse		10.9% (8;139) Worse
	San Joaquin Community Hospital				9.1% (22;112)	2.0% (6;367)	0.0% (0;37)	7.5% (17;246)	2.0% (10;578)	9.3% (62;579)	2.0% (6;326)	0.8% (1;155)	3.3% (11;461)
	Tehachapi Hospital								0.0% (0;3)				0.0% (0;39)
Kings	Adventist Medical Center								6.4% (2;39)	0.0% (0;7)	0.0% (0;19)	0.0% (0;4)	11.4% (4;53)
	Central Valley General Hospital							0.0% (0;3)	0.0% (0;61)	10.5% (1;26)	8.7% (1;21)		7.1% (5;107)
	Corcoran District Hospital												0.0% (0;20)
	Hanford Community Medical Center							13.6% (10;46) Worse	3.5% (9;298)	19.7% (19;73) Worse	3.0% (4;169)	7.6% (4;66)	8.3% (19;361) Worse
Lake	Saint Helena Hospital – Clearlake								0.0% (0;36)	32.7% (2;9)	3.7% (1;41)		3.0% (2;130)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Sutter Lakeside Hospital							0.0% (0;5)	0.0% (0;33)	4.1% (1;30)	0.0% (0;31)	1.8% (1;31)	0.0% (0;79)
Lassen	Banner Lassen Medical Center								5.9% (1;23)	19.5% (3;15)	0.0% (0;10)	0.0% (0;3)	28.3% (6;65) Worse
Los Angeles	Alhambra Hospital				55.9% (1;6)			7.9% (9;47)	1.7% (3;141)	7.4% (6;82)	0.8% (1;90)	2.0% (1;34)	1.7% (5;194) Better
	Antelope Valley Hospital			6.2% (1;8)	12.0% (11;72)	2.2% (7;298)	0.0% (0;39)	9.3% (33;393)	5.1% (20;524) Worse	10.4% (29;345)	3.7% (12;319)	4.4% (4;153)	7.4% (43;620) Worse
	Bellflower Medical Center							48.9% (2;7)	0.0% (0;59)	0.0% (0;8)	4.1% (2;55)	0.0% (0;8)	1.9% (2;102)
	Beverly Hospital				6.3% (1;8)	1.7% (3;121)	0.0% (0;14)	6.4% (12;158)	1.8% (7;397)	9.0% (11;130)	3.1% (5;130)	2.0% (1;61)	3.2% (11;282)
	Brotman Medical Center			8.2% (1;5)		1.9% (1;35)	0.0% (0;9)	7.5% (4;51)	1.1% (3;274)	8.6% (4;83)	0.0% (0;113)	3.0% (1;38)	4.3% (13;216)
	California Hospital Medical Center – Los Angeles		0.0% (0;3)		21.8% (7;25) Worse			8.7% (6;41)	1.4% (4;475)	11.3% (23;171)	2.7% (5;195)	4.8% (1;27)	6.2% (10;191)
	Catalina Island Medical Center												0.0% (0;3)
	Cedars Sinai Medical Center	9.0% (2;14)	3.4% (1;31)	0.0% (0;46)	3.3% (20;603) Better	2.6% (21;688)	1.3% (1;82)	8.6% (39;467)	3.6% (51;1,297)	7.6% (67;656) Better	2.3% (11;501)	1.6% (3;213)	3.7% (31;825)
	Centinela Hospital Medical Center			0.0% (0;6)	9.1% (13;45)	0.0% (0;194)	2.6% (1;22)	4.3% (14;252)	1.1% (12;1,187) Better	5.6% (28;406) Better	1.5% (7;369)	0.6% (1;73)	1.2% (5;332) Better
	Citrus Valley Medical Center – Inter Community Campus			0.0% (0;6)	3.3% (1;10)	3.5% (9;303)	0.0% (0;53)	7.7% (17;359)	1.3% (5;375) Better	8.9% (11;148)	2.8% (4;153)	0.0% (0;70)	3.5% (9;239)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Citrus Valley Medical Center – Queen of the Valley Campus			0.0% (0;3)	3.9% (2;54)		0.0% (0;23)	3.4% (4;58)	1.4% (5;364)	10.2% (21;184)	2.6% (5;183)	1.8% (1;70)	4.1% (8;204)
	City of Hope Helford Clinical Research Hospital	0.0% (0;9)	0.0% (0;14)		0.0% (0;106) Better				0.0% (0;3)	0.0% (0;7)	8.5% (2;26)	12.9% (1;3)	8.1% (7;102)
	Coast Plaza Doctors Hospital							6.6% (2;20)	1.2% (1;106)	7.8% (3;28)	5.7% (4;48)	0.0% (0;8)	7.7% (11;166)
	Community Hospital of Long Beach							4.1% (3;39)	12.2% (10;79) Worse	11.2% (4;48)	2.1% (1;61)	0.0% (0;30)	4.4% (10;142)
	Community and Mission Hospital of Huntington Park – Slauson							0.0% (0;10)	1.6% (1;115)	0.0% (0;37)	0.0% (0;57)		3.5% (2;101)
	Downey Regional Medical Center			0.0% (0;7)	9.6% (4;59)	2.3% (4;177)	0.0% (0;9)	6.4% (17;224)	3.2% (13;515)	8.2% (15;197)	2.1% (5;267)	0.0% (0;127)	3.7% (11;298)
	East Los Angeles Doctors Hospital							19.6% (7;24) Worse	0.7% (1;123)	8.3% (3;44)	0.0% (0;48)	0.0% (0;8)	3.0% (5;135)
	East Valley Hospital Medical Center							0.0% (0;16)	2.8% (2;50)	0.0% (0;12)	2.7% (1;25)	0.0% (0;4)	1.0% (2;121) Better
	Encino Hospital Medical Center							2.1% (1;17)	0.0% (0;56) Better	0.0% (0;18)	0.7% (1;58)	0.0% (0;27)	0.0% (0;33)
	Foothill Presbyterian Hospital – Johnston Memorial				0.0% (0;12)			0.0% (0;32)	3.7% (5;144)	6.0% (5;92)	0.0% (0;123)	0.0% (0;52)	7.7% (12;158) Worse
	Garfield Medical Center			0.0% (0;7)	4.8% (3;52)	1.8% (7;201)	0.0% (0;21)	5.5% (11;209)	3.5% (19;450)	4.7% (11;248) Better	0.6% (1;181)	0.0% (0;64)	3.6% (18;357)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Glendale Adventist Medical Center – Wilson Terrace	0.0% (0;3)	20.2% (1;5)	0.0% (0;15)	5.9% (14;127)	1.9% (15;673)	0.0% (0;22)	5.1% (22;400)	2.5% (22;537)	8.8% (48;403)	0.4% (1;259) Better	1.6% (3;149)	3.4% (21;400)
	Glendale Memorial Hospital and Medical Center			0.0% (0;14)	5.1% (2;23)	0.9% (3;300)	0.0% (0;28)	6.7% (17;275)	2.4% (14;521)	5.2% (10;175) Better	2.9% (6;201)	3.2% (2;90)	3.2% (13;299)
	Good Samaritan Hospital – Los Angeles		0.0% (0;4)	4.4% (1;11)	9.7% (5;45)	2.3% (11;391)	0.0% (0;11)	10.5% (30;296) Worse	4.0% (23;655)	9.3% (17;172)	2.0% (4;186)	2.2% (2;84)	4.7% (14;243)
	Greater El Monte Community Hospital							4.5% (2;18)	1.9% (1;66)	4.5% (1;31)	3.2% (2;43)		1.8% (2;89)
	Henry Mayo Newhall Memorial Hospital				12.3% (4;31)		0.0% (0;11)	6.3% (11;82)	2.7% (10;278)	13.9% (26;208)	1.3% (3;191)	0.9% (1;89)	2.6% (8;318)
	Hollywood Community Hospital							0.0% (0;21)	11.2% (3;61)	0.0% (0;10)	2.1% (1;49)	20.4% (1;6)	5.6% (5;137)
	Hollywood Presbyterian Medical Center			0.0% (0;3)	6.4% (2;14)		0.0% (0;14)	9.5% (13;86)	2.6% (12;400)	9.0% (15;134)	2.5% (5;182)	5.0% (4;65)	3.4% (11;251)
	Huntington Memorial Hospital	0.0% (0;3)	0.0% (0;7)	11.6% (1;11)	7.0% (7;190)	2.3% (8;345)	0.0% (0;39)	6.4% (19;294)	2.1% (10;509)	12.2% (50;409)	1.8% (5;279)	1.4% (2;179)	1.8% (6;417) Better
	Kaiser Foundation Hospital – Baldwin Park			0.0% (0;17)			0.0% (0;58)	4.6% (6;102)	2.0% (7;439)	8.4% (13;238)	1.7% (4;325)	3.4% (3;112)	3.2% (12;407)
	Kaiser Foundation Hospital – Downey	0.0% (0;3)	0.0% (0;4)	0.0% (0;24)			0.0% (0;26)	6.5% (12;142)	1.0% (5;613) Better	12.2% (20;254)	2.5% (6;344)	4.9% (3;84)	4.0% (12;360)
	Kaiser Foundation Hospital – South Bay*		0.0% (0;10)	2.9% (1;20)			0.0% (0;56)	3.9% (5;93)	3.0% (12;549)	7.2% (11;225)	2.6% (7;325)	5.5% (3;74)	4.0% (11;318)
	Kaiser Foundation Hospital – Panorama City	0.0% (0;5)	0.0% (0;5)	0.0% (0;6)		0.0% (0;26)	0.0% (0;26)	3.7% (5;107)	2.1% (8;398)	8.0% (14;210)	2.6% (7;247)	2.4% (3;119)	4.8% (27;472)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Kaiser Foundation Hospital – Sunset	0.0% (0;7)	0.0% (0;13)	0.0% (0;18)	6.1% (17;432)	1.6% (11;1,041)	0.0% (0;14)	6.0% (43;1,103)	3.7% (22;619)	8.1% (44;476)	2.0% (6;276)	3.0% (2;48)	3.9% (13;303)
	Kaiser Foundation Hospital – West Los Angeles	0.0% (0;3)		0.0% (0;12)			0.0% (0;18)	3.0% (3;98)	2.9% (12;452)	6.9% (11;228)	2.1% (6;328)	0.0% (0;40)	4.3% (14;282)
	Kaiser Foundation Hospital – Woodland Hills			0.0% (0;19)	5.2% (4;68)		1.1% (1;75)	5.1% (8;113)	4.8% (24;364) Worse	9.3% (29;288)	1.7% (4;285)	2.8% (6;179)	4.5% (16;268)
	Lakewood Regional Medical Center			21.1% (1;5)	8.8% (3;28)	2.1% (7;337)	0.0% (0;20)	9.4% (23;279)	4.5% (16;450)	10.1% (17;179)	2.6% (4;167)	3.0% (3;100)	4.6% (8;214)
	Lancaster Community Hospital			45.7% (1;5)	9.9% (4;18)	5.4% (4;159)	0.0% (0;8)	11.4% (12;125)	2.7% (5;261)	12.3% (14;119)	2.6% (3;123)	2.2% (1;68)	3.6% (8;223)
	Long Beach Memorial Medical Center	0.0% (0;3)		0.0% (0;29)	10.6% (15;156)	1.5% (7;457)	1.3% (1;90)	7.0% (19;347)	3.0% (15;696)	10.8% (60;560)	3.4% (10;350)	0.8% (1;141)	4.6% (20;457)
	Los Angeles Community Hospital							6.8% (2;22)	4.9% (4;150)	3.3% (1;65)	0.0% (0;57)	0.0% (0;9)	5.5% (8;122)
	Los Angeles County/Harbor – UCLA Medical Center		0.0% (0;4)	0.0% (0;15)	7.5% (5;91)	2.5% (4;368)	0.0% (0;7)	6.4% (9;295)	2.6% (8;560)	10.2% (15;221)	3.1% (5;231)	0.0% (0;30)	5.3% (11;293)
	Los Angeles County/Olive View – UCLA Medical Center		0.0% (0;4)				0.0% (0;7)	6.8% (3;44)	1.4% (3;383)	7.4% (4;109)	1.5% (1;144)		4.5% (8;318)
	Los Angeles County/University of Southern California Medical Center		10.3% (1;14)	0.0% (0;7)	15.7% (18;225) Worse	2.3% (7;417)	0.0% (0;20)	12.5% (18;295) Worse	3.6% (10;753)	18.5% (45;221) Worse	3.2% (7;326)	5.4% (2;32)	5.0% (10;471)
	Los Angeles Metropolitan Medical Center							0.0% (0;4)	5.3% (4;111)	0.0% (0;18)	3.8% (1;34)	0.0% (0;6)	3.6% (5;104)
	Marina Del Rey Hospital							4.7% (4;52)	1.8% (4;179)	4.0% (2;69)	0.0% (0;83)	1.6% (1;43)	3.9% (7;124)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Memorial Hospital of Gardena							6.2% (4;38)	1.3% (3;269)	3.2% (3;118) Better	1.5% (4;146)	0.0% (0;18)	3.1% (5;133)
	Methodist Hospital of Southern California			6.6% (1;4)	6.4% (2;28)	1.7% (6;327)	0.0% (0;42)	6.7% (26;341)	2.9% (15;511)	9.3% (33;360)	0.4% (1;241)	1.6% (3;190)	3.0% (14;395)
	Mission Community Hospital – Panorama Campus							6.6% (7;43)	1.4% (3;139)	9.3% (2;23)	0.0% (0;83)	3.0% (1;27)	1.3% (1;80)
	Monterey Park Hospital							2.6% (1;23)	1.5% (2;128)	5.1% (1;33)	2.6% (2;88)	0.0% (0;21)	3.7% (4;109)
	Motion Picture and Television Hospital								0.0% (0;10)	0.0% (0;3)	0.0% (0;17)		30.1% (3;15) Worse
	Northridge Hospital Medical Center		15.1% (1;5)	0.0% (0;7)	10.6% (9;45)	1.7% (7;338)	0.0% (0;38)	5.8% (19;293)	1.2% (6;408) Better	9.9% (37;304)	2.3% (6;215)	2.2% (4;147)	2.7% (12;310)
	Norwalk Community Hospital							3.0% (1;22)	0.6% (1;98)	8.4% (2;19)	4.8% (2;38)	0.0% (0;6)	4.6% (2;61)
	Olympia Medical Center						0.0% (0;3)	3.8% (4;58)	3.2% (9;272)	6.0% (3;85)	1.7% (3;138)	0.0% (0;45)	2.8% (11;224)
	Pacific Alliance Medical Center, Inc.				0.0% (0;6)			0.0% (0;18)	3.1% (7;176)	7.0% (3;79)	1.5% (2;82)	4.5% (1;17)	6.6% (8;119)
	Pacific Hospital of Long Beach							0.0% (0;11)	2.2% (2;82)	9.0% (1;21)	0.9% (1;80)	3.4% (1;15)	2.6% (4;105)
	Pacifica Hospital of the Valley							14.4% (5;13)	2.7% (2;79)	11.6% (4;34)	13.6% (6;61) Worse	30.9% (1;9)	1.4% (2;98)
	Pomona Valley Hospital Medical Center		0.0% (0;3)	0.0% (0;9)	11.2% (6;82)	1.3% (5;240)	7.4% (1;10)	4.8% (14;276)	2.3% (16;570)	9.9% (29;346)	2.2% (7;318)	2.3% (3;126)	3.9% (12;286)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Presbyterian Intercommunity Hospital			0.0% (0;8)	2.6% (1;40)	0.3% (1;313) Better	0.0% (0;37)	4.2% (26;552) Better	2.3% (13;549)	5.5% (24;389) Better	1.0% (3;320)	1.3% (2;181)	1.6% (6;363) Better
	Providence Holy Cross Medical Center	0.0% (0;3)		0.0% (0;9)	13.3% (6;52)	2.7% (12;379)	0.0% (0;46)	8.2% (14;264)	1.9% (12;473)	9.0% (24;280)	1.1% (3;235)	0.8% (1;117)	3.5% (17;386)
	Providence Little Company of Mary Medical Center – San Pedro				6.9% (3;15)		8.4% (1;16)	9.8% (4;24)	2.6% (4;180)	8.4% (19;145)	1.8% (2;120)	4.4% (3;69)	2.1% (3;145)
	Providence Little Company of Mary Medical Center – Torrance			0.0% (0;9)	7.7% (3;37)	1.7% (6;413)	2.3% (1;25)	7.4% (22;308)	2.7% (15;581)	8.4% (35;336)	1.3% (4;326)	1.5% (3;172)	3.6% (17;410)
	Providence Saint Joseph Medical Center	0.0% (0;3)		0.0% (0;11)	12.6% (13;123) Worse	3.5% (12;351)	1.3% (1;58)	7.3% (13;269)	3.2% (22;545)	11.8% (48;403)	3.8% (10;293)	2.4% (4;194)	3.4% (13;288)
	Providence Tarzana Medical Center			1.6% (1;39)	9.1% (1;10)	1.8% (5;270)	0.9% (1;103)	5.1% (9;159)	2.4% (13;371)	8.6% (13;167)	1.1% (2;166)	0.7% (1;112)	2.1% (11;329) Better
	Ronald Reagan UCLA Medical Center	0.0% (0;15)	0.0% (0;55)	0.0% (0;66)	4.6% (28;674) Better	3.1% (11;190)	0.0% (0;42)	6.2% (12;185)	3.6% (17;386)	9.2% (70;498)	2.0% (4;159)	7.1% (3;43)	5.2% (13;254)
	Saint Francis Medical Center			0.0% (0;6)	4.4% (1;25)	3.6% (7;146)	0.0% (0;13)	9.5% (19;169)	2.5% (15;575)	8.9% (23;178)	2.4% (6;215)	3.0% (2;45)	2.7% (7;217)
	Saint John’s Health Center		9.7% (1;7)	0.0% (0;8)	1.3% (2;131) Better	1.3% (3;335)	0.0% (0;42)	5.4% (9;170)	3.0% (11;308)	5.5% (8;142) Better	1.4% (4;214)	1.3% (2;144)	3.3% (11;293)
	Saint Mary Medical Center			0.0% (0;3)	0.0% (0;17)	2.3% (3;180)	0.0% (0;18)	6.8% (14;205)	1.0% (3;297) Better	5.7% (8;133)	1.1% (3;185)	0.0% (0;60)	4.6% (21;319)
	Saint Vincent Medical Center		0.0% (0;6)	0.0% (0;12)	1.8% (1;236)	1.4% (3;115)	0.0% (0;14)	7.5% (11;133)	3.4% (12;423)	9.4% (8;128)	0.7% (1;142)	0.0% (0;47)	4.3% (10;282)
	San Dimas Community Hospital							6.3% (2;16)	4.7% (4;72)	7.8% (6;67)	1.6% (1;50)	0.0% (0;35)	1.1% (1;73)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	San Gabriel Valley Medical Center				6.5% (1;17)			8.4% (9;47)	2.4% (8;270)	8.9% (9;122)	2.1% (3;145)	4.3% (3;66)	2.7% (13;328)
	Santa Monica – UCLA Medical Center and Orthopedic Hospital				8.9% (1;14)	2.1% (3;103)	0.0% (0;4)	4.1% (7;134)	2.3% (10;331)	12.7% (13;128)	1.5% (3;223)	2.6% (4;133)	3.6% (12;350)
	Sherman Oaks Hospital							3.5% (10;129) Better	0.5% (1;96) Better	7.8% (6;59)	0.0% (0;75) Better	0.0% (0;37)	2.4% (3;87)
	Silver Lake Medical Center – Downtown Campus							0.0% (0;5)	0.0% (0;92)	0.0% (0;11)	3.9% (1;43)	0.0% (0;5)	0.0% (0;49)
	Temple Community Hospital							8.9% (4;39)	3.7% (3;140)	4.6% (1;33)	2.1% (1;51)		3.1% (2;83)
	Torrance Memorial Medical Center			0.0% (0;32)	7.1% (5;57)	1.8% (15;717)	0.0% (0;80)	6.4% (31;439)	3.8% (24;723)	9.4% (52;501)	2.3% (9;425)	2.6% (6;242)	4.0% (23;532)
	Tri-City Regional Medical Center – Hawaiian Gardens							5.8% (3;18)	2.9% (4;122)	11.7% (1;17)	3.3% (4;89)	0.0% (0;14)	0.0% (0;116)
	University of Southern California Kenneth Norris, Jr. Cancer Hospital										0.0% (0;4)		
	University of Southern California University Hospital*	0.0% (0;49)	0.0% (0;40)	5.9% (2;40)	3.9% (6;273)	3.0% (7;89)	0.0% (0;19)	15.2% (11;57) Worse	3.4% (4;127)	7.4% (10;127)	0.0% (0;45)	19.8% (1;8)	9.5% (8;62) Worse
	Valley Presbyterian Hospital			0.0% (0;3)	5.1% (1;18)	1.8% (7;254)	0.0% (0;16)	7.0% (25;293)	2.6% (11;355)	13.1% (17;128)	3.1% (4;118)	3.1% (2;69)	5.8% (17;216)
	Verdugo Hills Hospital				0.0% (0;5)			10.5% (7;42)	3.9% (7;188)	4.5% (2;69)	3.3% (3;109)	1.7% (1;71)	5.2% (8;168)
	West Hills Hospital and Medical Center				15.0% (3;22)	3.2% (3;143)	0.0% (0;24)	7.7% (14;167)	5.5% (12;180)	10.7% (12;141)	3.8% (7;167)	5.3% (5;132)	4.3% (18;283)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	White Memorial Medical Center			0.0% (0;5)	16.7% (7;25) Worse	1.5% (9;314)	0.0% (0;16)	5.2% (13;225)	1.4% (7;458) Better	9.5% (44;333)	0.3% (1;237) Better	0.9% (1;57)	0.7% (2;270) Better
	Whittier Hospital Medical Center							3.0% (4;77)	2.5% (5;175)	3.6% (1;68)	0.5% (1;175)	4.7% (2;38)	3.4% (4;144)
Madera	Madera Community Hospital							14.2% (10;57) Worse	7.3% (7;126)	6.9% (3;35)	2.5% (2;70)	0.0% (0;13)	9.6% (17;218) Worse
Marin	Kaiser Foundation Hospital – San Rafael			0.0% (0;5)		3.0% (2;24)	0.0% (0;14)	6.2% (7;97)	5.6% (10;186)	12.4% (12;151)	4.3% (4;118)	4.2% (3;91)	5.5% (9;207)
	Marin General Hospital			0.0% (0;13)	8.4% (6;24)	1.8% (2;187)	0.0% (0;36)	3.4% (5;177)	5.0% (10;189)	11.3% (24;190)	2.5% (4;173)	2.2% (3;116)	4.8% (14;286)
	Novato Community Hospital							0.0% (0;11)	1.1% (1;62)	0.0% (0;18)	0.0% (0;32)	4.1% (1;30)	1.8% (1;43)
Mariposa	John C. Fremont Healthcare District							0.0% (0;4)	21.7% (1;5)	42.9% (1;5)			12.0% (1;18)
Mendocino	Frank R. Howard Memorial Hospital								0.0% (0;28)	19.1% (2;19)	10.8% (1;14)	7.9% (1;25)	7.4% (2;47)
	Mendocino Coast District Hospital							4.4% (1;14)	0.0% (0;25)	8.7% (1;20)	0.0% (0;35)	0.0% (0;14)	12.3% (4;56)
	Ukiah Valley Medical Center							10.2% (2;6)	3.0% (6;135)	4.7% (3;63)	0.0% (0;47)	13.9% (4;37) Worse	3.2% (6;160)
Merced	Memorial Hospital Los Banos							7.3% (1;7)	11.6% (4;40) Worse	0.0% (0;18)	0.0% (0;7)		5.2% (3;60)
	Mercy Medical Center – Merced							6.7% (11;90)	3.2% (16;351)	12.0% (18;201)	2.3% (6;241)	0.0% (0;66)	4.8% (21;372)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
Modoc	Modoc Medical Center								0.0% (0;14)	0.0% (0;3)			9.0% (1;25)
Mono	Mammoth Hospital										0.0% (0;4)	0.0% (0;6)	0.0% (0;7)
Monterey	Community Hospital Monterey Peninsula			6.0% (1;4)	2.7% (1;25)	1.2% (2;170)	0.0% (0;48)	9.7% (12;139)	3.2% (7;239)	12.5% (22;196)	2.8% (8;228)	3.3% (3;122)	7.4% (20;272) Worse
	George L. Mee Memorial Hospital								0.0% (0;39)	0.0% (0;7)	0.0% (0;13)	0.0% (0;4)	7.1% (2;35)
	Natividad Medical Center							10.2% (3;12)	2.7% (2;87)	24.4% (2;32)	4.3% (2;71)	0.0% (0;5)	7.3% (6;147)
	Salinas Valley Memorial Hospital			0.0% (0;4)	4.6% (3;48)	2.5% (9;235)	0.0% (0;29)	5.0% (11;240)	4.5% (17;338)	10.5% (21;192)	3.0% (5;162)	1.5% (1;93)	4.7% (13;227)
Napa	Queen of the Valley Hospital – Napa			0.0% (0;13)		2.6% (6;172)	0.0% (0;20)	8.6% (20;201)	2.1% (7;265)	13.6% (13;118)	2.0% (4;148)	4.1% (3;76)	4.2% (10;215)
	Saint Helena Hospital			0.0% (0;10)		1.3% (2;282)	1.6% (1;64)	7.0% (6;86)	2.1% (3;140)	10.1% (3;61)	0.0% (0;87)	3.4% (1;29)	1.0% (1;83)
Nevada	Sierra Nevada Memorial Hospital						0.0% (0;12)	3.4% (2;41)	3.2% (7;173)	14.1% (12;109)	2.3% (2;92)	4.1% (4;88)	5.9% (10;199)
	Tahoe Forest Hospital						0.0% (0;3)		17.0% (2;19)	12.7% (1;14)	0.0% (0;13)	0.0% (0;25)	0.0% (0;43)
Orange	AHMC Anaheim Regional Medical Center			0.0% (0;17)	0.0% (0;23)	2.0% (4;229)	0.8% (1;56)	7.0% (18;273)	1.2% (5;535) Better	8.1% (13;202)	1.5% (3;241)	2.6% (2;96)	1.7% (5;325) Better
	Anaheim General Hospital								0.0% (0;20)	0.0% (0;3)	0.0% (0;9)		3.5% (1;15)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Chapman Medical Center							15.3% (1;3)	7.5% (2;35)	0.0% (0;14)	4.1% (1;26)	0.0% (0;15)	6.5% (3;59)
	Coastal Communities Hospital							6.2% (1;13)	1.6% (1;80)	0.0% (0;22)	1.0% (1;71)	0.0% (0;10)	7.8% (7;127)
	College Hospital Costa Mesa												0.0% (0;8)
	Fountain Valley Regional Hospital and Medical Center – Euclid		0.0% (0;8)	0.0% (0;22)	8.2% (6;55)	1.1% (4;257)	0.0% (0;25)	6.6% (20;310)	2.4% (11;487)	9.1% (27;297)	2.3% (5;240)	1.9% (2;88)	1.4% (5;347) Better
	Garden Grove Hospital and Medical Center				3.9% (1;14)			8.1% (10;64)	2.3% (3;106)	9.5% (9;95)	3.8% (5;96)	3.0% (1;40)	0.7% (1;106) Better
	Hoag Memorial Hospital Presbyterian	0.0% (0;8)		1.6% (1;35)	5.9% (22;340)	1.4% (7;380)	0.0% (0;92)	5.5% (17;354)	3.7% (25;552)	7.4% (46;547) Better	2.1% (8;412)	2.5% (5;275)	4.6% (27;589)
	Hoag Orthopedic Institute											0.0% (0;10)	
	Huntington Beach Hospital							4.8% (4;32)	0.4% (1;130) Better	0.0% (0;36) Better	1.1% (2;75)	0.0% (0;20)	0.0% (0;80) Better
	Kaiser Foundation Hospital – Anaheim		19.6% (1;7)	4.5% (1;33)	8.2% (10;200)	0.0% (0;13)	0.0% (0;57)	8.8% (20;165)	3.4% (17;491)	8.0% (36;470)	2.2% (7;378)	3.0% (5;185)	2.9% (16;548)
	La Palma Intercommunity Hospital							5.0% (5;50)	0.8% (1;130)	11.0% (3;45)	0.0% (0;75)	0.0% (0;35)	1.5% (2;108)
	Los Alamitos Medical Center			0.0% (0;11)	15.9% (7;27) Worse	4.6% (4;84)	0.0% (0;50)	12.2% (17;135) Worse	3.9% (11;282)	10.5% (32;326)	4.4% (6;186)	1.1% (1;139)	7.0% (21;305) Worse
	Mission Hospital Laguna Beach				0.0% (0;10)			7.6% (1;8)	2.1% (1;37)	6.5% (1;19)	0.0% (0;48)	0.0% (0;43)	3.3% (2;67)
	Mission Hospital Regional Medical Center			0.0% (0;8)	4.3% (8;143)	2.0% (9;460)	0.0% (0;12)	7.6% (16;258)	3.3% (16;363)	8.4% (43;475)	2.4% (6;254)	0.8% (1;132)	2.9% (8;278)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Orange Coast Memorial Medical Center		0.0% (0;5)		0.0% (0;23)	1.5% (1;96)	1.7% (1;20)	7.1% (14;143)	3.0% (9;334)	5.1% (6;154) Better	0.5% (1;196)	1.0% (1;132)	4.2% (12;238)
	Placentia Linda Hospital							6.2% (3;32)	3.0% (4;168)	9.7% (3;57)	2.8% (2;110)	1.4% (1;86)	3.4% (5;185)
	Saddleback Memorial Medical Center			0.0% (0;19)	10.6% (2;42)	0.9% (4;378)	0.0% (0;42)	4.1% (19;313) Better	2.4% (20;595)	11.6% (35;319)	0.9% (3;280)	1.7% (6;281)	3.4% (18;430)
	Saint Joseph Hospital – Orange	0.0% (0;3)	0.0% (0;3)	0.0% (0;57)	3.9% (9;166)	1.7% (4;223)	0.0% (0;88)	5.8% (12;214)	3.0% (16;384)	4.8% (19;330) Better	0.9% (3;237)	1.7% (4;179)	2.5% (9;276)
	Saint Jude Medical Center			0.0% (0;27)	6.5% (8;68)	0.9% (4;370)	0.0% (0;58)	4.1% (15;371) Better	1.9% (12;578)	9.4% (37;362)	1.5% (4;281)	3.4% (6;211)	4.6% (20;382)
	UC Irvine Medical Center	0.0% (0;17)	2.9% (1;33)	0.0% (0;17)	6.4% (15;156)	1.2% (2;130)	0.0% (0;26)	4.1% (4;95)	0.6% (1;223)	11.0% (54;269)	1.7% (3;135)	2.3% (1;21)	2.7% (4;166)
	West Anaheim Medical Center					1.0% (2;114)	0.0% (0;3)	3.9% (6;124)	2.1% (8;249)	1.2% (1;83) Better	3.0% (8;125)	0.0% (0;47)	2.1% (10;268) Better
	Western Medical Center – Anaheim					0.0% (0;97)	0.0% (0;3)	4.7% (4;98)	6.0% (3;45)	0.0% (0;16)	0.0% (0;41)	0.0% (0;12)	5.4% (3;52)
	Western Medical Center – Santa Ana				8.4% (4;55)	1.2% (4;309)	0.0% (0;22)	7.0% (15;253)	4.9% (8;182)	12.6% (24;198)	1.9% (2;103)	0.0% (0;36)	3.5% (5;146)
Placer	Sutter Auburn Faith Hospital			0.0% (0;6)			0.0% (0;13)	4.1% (2;25)	1.6% (2;109)	8.1% (8;90)	1.8% (2;66)	2.4% (2;85)	4.5% (7;147)
	Sutter Roseville Medical Center		37.6% (1;3)	0.0% (0;11)	6.9% (9;62)	2.1% (5;205)	0.0% (0;35)	6.7% (16;236)	0.9% (4;316) Better	9.7% (36;302)	1.7% (6;291)	3.1% (7;173)	1.5% (6;364) Better
Plumas	Eastern Plumas Hospital – Portola Campus								11.8% (2;20)	0.0% (0;3)	0.0% (0;8)		0.0% (0;22)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Plumas District Hospital								0.0% (0;10)	60.8% (1;5)	0.0% (0;9)		7.0% (1;29)
	Seneca Healthcare District								0.0% (0;11)				0.0% (0;32)
Riverside	Corona Regional Medical Center – Main			9.5% (1;7)			0.0% (0;16)	9.3% (10;66)	4.3% (14;311)	12.0% (10;109)	4.0% (6;139)	2.3% (2;75)	6.6% (17;228)
	Desert Regional Medical Center			4.0% (1;18)	5.8% (7;89)	1.6% (9;464)	0.0% (0;52)	5.5% (16;352)	2.9% (10;398)	7.8% (36;382)	2.6% (6;245)	2.1% (3;160)	4.0% (11;345)
	Eisenhower Medical Center			0.0% (0;24)	9.4% (8;68)	2.9% (11;610)	0.0% (0;93)	7.2% (27;426)	2.5% (16;607)	8.8% (33;435)	2.2% (9;418)	2.0% (6;284)	4.0% (29;680)
	Hemet Valley Medical Center			0.0% (0;9)			0.0% (0;21)	9.0% (22;146)	2.7% (12;455)	10.0% (19;224)	4.6% (12;233) Worse	4.1% (6;169)	5.1% (17;322)
	John F. Kennedy Memorial Hospital					2.8% (1;35)	0.0% (0;7)	7.3% (5;86)	2.5% (3;149)	4.6% (1;43)	4.0% (2;79)	1.7% (1;72)	3.6% (6;194)
	Kaiser Foundation Hospital – Moreno Valley						0.0% (0;5)	11.6% (5;18)	2.5% (3;116)	8.2% (2;43)	2.5% (2;83)	3.7% (2;30)	2.1% (2;110)
	Kaiser Foundation Hospital – Riverside	0.0% (0;11)	10.1% (1;9)	0.0% (0;8)			0.0% (0;52)	6.3% (10;130)	3.9% (13;344)	8.6% (13;188)	2.5% (5;243)	3.7% (3;112)	3.1% (9;303)
	Menifee Valley Medical Center						0.0% (0;16)	9.5% (15;77)	4.1% (7;187)	10.4% (7;76)	1.3% (1;114)	5.7% (3;93)	6.9% (15;138)
	Palo Verde Hospital							9.0% (1;11)	30.6% (4;29) Worse	0.0% (0;6)	0.0% (0;37)		17.1% (3;61)
	Parkview Community Hospital Medical Center						0.0% (0;11)	10.9% (7;34)	2.4% (5;231)	9.5% (5;76)	2.0% (3;132)	2.8% (2;47)	4.2% (10;250)
	Riverside Community Hospital			5.9% (1;19)	9.3% (7;64)	2.0% (25;817)	0.0% (0;43)	8.7% (45;709)	2.6% (15;658)	9.5% (38;374)	2.8% (9;361)	1.4% (2;154)	4.5% (16;455)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia	
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.														
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)	
	Riverside County Regional Medical Center	12.5% (1;4)			11.2% (4;41)		24.8% (1;17) Worse	6.3% (4;66)	2.9% (6;418)	15.3% (25;196) Worse	4.6% (6;146)	1.8% (1;46)	6.4% (13;282)	
	San Geronio Memorial Hospital							2.7% (3;62)	1.5% (3;170)	9.8% (3;51)	2.4% (2;68)	3.3% (1;40)	5.9% (13;210)	
	Southwest Healthcare System – Murrieta			0.0% (0;15)	14.1% (10;82) Worse		0.0% (0;75)	4.7% (6;97)	1.6% (6;462)	10.9% (27;306)	1.4% (2;244)	1.6% (2;170)	3.7% (14;422)	
Sacramento	Kaiser Foundation Hospital – Sacramento/Roseville-Morse*	0.0% (0;6)	0.0% (0;16)	6.3% (2;30)	5.5% (20;310)	2.5% (3;99)	0.0% (0;82)	6.9% (31;409)	5.5% (43;848) Worse	10.4% (89;825)	2.6% (12;528)	4.7% (15;393) Worse	5.9% (34;670) Worse	
	Kaiser Foundation Hospital – South Sacramento	35.2% (1;5)	0.0% (0;5)	0.0% (0;20)	79.5% (1;4)		0.0% (0;31)	5.3% (15;162)	3.2% (13;397)	12.1% (25;291)	3.1% (4;202)	2.5% (3;118)	4.2% (12;298)	
	Mercy General Hospital			3.2% (1;23)	5.9% (13;141)	2.7% (29;1,646)	0.0% (0;83)	5.5% (36;1,099)	1.3% (6;498) Better	7.6% (36;358) Better	3.7% (9;167)	2.5% (3;98)	3.1% (10;300)	
	Mercy Hospital – Folsom							11.8% (5;18)	3.5% (3;90)	9.2% (4;57)	4.7% (3;92)	0.0% (0;54)	4.2% (4;139)	
	Mercy San Juan Hospital	0.0% (0;5)		0.0% (0;39)	9.0% (12;84)	2.1% (6;260)	0.0% (0;99)	7.6% (22;310)	3.4% (21;471)	13.0% (62;447) Worse	3.0% (9;280)	2.8% (6;185)	4.3% (20;413)	
	Methodist Hospital of Sacramento		0.0% (0;3)					7.9% (7;37)	3.3% (7;241)	9.9% (12;171)	2.9% (6;197)	3.4% (3;104)	3.3% (6;239)	
	Sutter General Hospital	29.6% (1;5)	0.0% (0;6)	0.0% (0;7)	4.8% (5;130)			0.0% (0;41)	11.1% (7;27)	3.2% (8;234)	8.8% (25;243)	1.5% (4;239)	1.6% (4;163)	3.3% (9;286)
	Sutter Memorial Hospital			0.0% (0;41)	3.0% (1;98)	2.2% (9;373)	0.0% (0;116)	6.2% (24;472)	3.7% (14;343)	5.8% (4;77)	1.1% (1;63)		2.8% (5;179)	
	UC Davis Medical Center	0.0% (0;12)	6.8% (1;14)	3.0% (1;19)	9.3% (14;263)	2.4% (17;551)	0.0% (0;30)	5.6% (20;408)	2.6% (12;569)	13.0% (45;328) Worse	2.2% (7;287)	5.4% (3;81)	3.8% (14;411)	

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
San Benito	Hazel Hawkins Memorial Hospital							0.0% (0;8)	0.0% (0;73)	10.2% (2;29)	0.0% (0;45)	0.0% (0;14)	8.1% (6;108)
San Bernardino	Arrowhead Regional Medical Center		27.6% (1;5)		16.5% (12;95) Worse	5.2% (2;89)	0.0% (0;12)	12.5% (11;117) Worse	7.1% (13;514) Worse	14.0% (60;426) Worse	4.9% (4;138)	12.0% (2;28)	6.3% (15;306)
	Barstow Community Hospital							0.0% (0;19)	0.0% (0;150) Better	0.0% (0;46)	0.0% (0;31)	0.0% (0;6)	0.0% (0;112) Better
	Bear Valley Community Hospital							0.0% (0;3)	0.0% (0;21)	0.0% (0;3)	0.0% (0;7)		14.6% (2;38)
	Chino Valley Medical Center							3.5% (3;36)	0.4% (1;170) Better	3.1% (3;86) Better	0.9% (1;91)	0.0% (0;27)	1.6% (2;93)
	Colorado River Medical Center								28.2% (1;8)	0.0% (0;3)	0.0% (0;12)		0.0% (0;17)
	Community Hospital of San Bernardino						0.0% (0;3)	5.9% (4;40)	3.1% (4;197)	2.2% (1;68)	2.2% (2;90)	0.0% (0;12)	5.4% (9;174)
	Desert Valley Hospital			0.0% (0;14)		1.5% (2;90)		8.3% (11;73)	0.8% (3;302) Better	5.6% (7;138)	0.5% (1;131)	0.0% (0;57) Better	1.1% (5;279) Better
	Hi-Desert Medical Center						0.0% (0;5)	25.4% (6;13) Worse	5.6% (6;153)	13.4% (2;51)	0.0% (0;40)	0.0% (0;49)	8.2% (9;144)
	Kaiser Foundation Hospital – Fontana	0.0% (0;13)	10.4% (1;8)	2.6% (1;23)	5.0% (5;152)		0.0% (0;40)	5.2% (16;235)	2.6% (20;763)	6.6% (31;521) Better	2.5% (9;451)	1.4% (2;188)	2.3% (17;619) Better
	Loma Linda University Medical Center	18.6% (2;10)	0.0% (0;3)	0.0% (0;26)	9.9% (30;350) Worse	3.1% (9;271)	0.0% (0;25)	7.2% (27;481)	4.3% (19;533)	11.1% (86;522)	0.9% (2;214)	0.0% (0;56)	4.9% (12;282)
	Montclair Hospital Medical Center							7.4% (2;14)	0.7% (1;122)	11.3% (2;33)	0.0% (0;53)	0.0% (0;15)	3.1% (7;139)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Mountains Community Hospital								0.0% (0;7)	0.0% (0;3)	0.0% (0;3)		13.0% (1;33)
	Redlands Community Hospital			0.0% (0;15)	0.0% (0;3)		0.0% (0;43)	11.3% (24;122) Worse	3.5% (10;251)	10.9% (11;118)	3.1% (6;196)	1.1% (2;186)	4.2% (22;464)
	Saint Bernadine Medical Center			0.0% (0;19)	8.3% (4;42)	1.6% (16;1,672)	0.0% (0;43)	5.3% (33;1,008)	1.4% (6;561) Better	8.7% (25;273)	1.8% (3;190)	1.4% (1;80)	2.1% (7;340)
	Saint Mary Regional Medical Center			0.0% (0;17)	0.0% (0;12)	2.9% (10;470)	0.0% (0;32)	6.2% (26;526)	2.5% (10;527)	5.7% (5;239)	2.2% (4;259)	1.5% (1;112)	4.6% (15;336)
	San Antonio Community Hospital			0.0% (0;11)	6.3% (5;60)	1.1% (3;248)	0.0% (0;43)	5.6% (26;468)	2.3% (16;528)	8.3% (30;322)	3.3% (8;220)	1.4% (2;150)	3.2% (18;437)
	Victor Valley Community Hospital							4.1% (1;29)	2.2% (3;162)	19.6% (4;52)	0.0% (0;75)	0.0% (0;29)	3.9% (5;159)
San Diego	Alvarado Hospital	0.0% (0;3)		0.0% (0;8)	10.6% (2;13)	1.7% (6;483)	0.0% (0;24)	5.6% (8;145)	3.8% (10;261)	8.7% (12;158)	3.4% (5;148)	3.9% (2;71)	4.3% (12;275)
	Fallbrook Hospital District							4.0% (1;14)	5.3% (4;64)	6.7% (1;19)	3.2% (1;38)	6.9% (2;34)	3.6% (2;71)
	Grossmont Hospital			0.0% (0;22)	5.3% (6;112)	1.7% (12;568)	0.0% (0;104)	6.4% (32;488)	4.7% (32;760) Worse	12.0% (65;574)	1.8% (7;413)	4.2% (8;217)	6.0% (30;637)
	Kaiser Foundation Hospital – San Diego	0.0% (0;6)	0.0% (0;8)	0.0% (0;39)	9.7% (7;212)		1.3% (1;85)	6.9% (22;192)	3.2% (22;817)	11.9% (38;486)	4.2% (14;439) Worse	5.0% (14;344) Worse	5.2% (36;930)
	Palomar Medical Center			9.3% (1;20)	5.1% (4;63)	2.7% (12;336)	0.0% (0;38)	9.2% (28;345)	3.1% (17;508)	11.3% (56;469)	1.3% (3;325)	1.4% (2;170)	3.1% (13;425)
	Paradise Valley Hospital				0.0% (0;7)	0.0% (0;10)	0.0% (0;7)	9.7% (8;45)	0.3% (1;261) Better	3.4% (3;98) Better	2.6% (4;100)	0.0% (0;19)	1.1% (3;197) Better
	Pomerado Hospital			0.0% (0;7)	0.0% (0;3)		0.0% (0;7)	8.0% (8;53)	3.2% (12;237)	9.6% (10;126)	1.6% (2;161)	0.0% (0;93)	3.9% (10;184)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Scripps Green Hospital		0.0% (0;9)	0.0% (0;28)	9.5% (6;93)	0.8% (6;1,405) Better	0.0% (0;51)	5.2% (9;234)	3.3% (10;321)	5.8% (10;175) Better	0.5% (1;179)	0.0% (0;100)	2.6% (6;211)
	Scripps Memorial Hospital – Encinitas			0.0% (0;11)	6.0% (3;27)	0.0% (0;129)	0.0% (0;34)	5.8% (9;164)	1.7% (5;282)	8.7% (21;240)	2.9% (6;174)	0.0% (0;171) Better	4.2% (11;220)
	Scripps Memorial Hospital – La Jolla			0.0% (0;11)	7.9% (13;179)	2.4% (19;1,307)	2.6% (1;36)	7.6% (25;505)	2.9% (9;273)	13.7% (44;273) Worse	1.5% (2;143)	0.7% (1;138)	2.0% (4;201)
	Scripps Mercy Hospital		0.0% (0;5)	0.0% (0;10)	6.5% (5;57)	1.2% (8;461)	0.0% (0;53)	6.2% (31;405)	2.0% (17;859)	10.8% (80;605)	2.4% (12;486)	1.8% (4;204)	4.4% (26;591)
	Sharp Chula Vista Medical Center		0.0% (0;4)	0.0% (0;15)	9.3% (3;36)	2.0% (9;392)	0.0% (0;31)	7.3% (33;349)	2.1% (12;598)	8.3% (12;193)	3.0% (8;256)	4.6% (3;91)	3.4% (12;373)
	Sharp Coronado Hospital and Healthcare Center							0.0% (0;10)	0.0% (0;59)	5.5% (1;25)	0.0% (0;37)	0.0% (0;31)	2.8% (1;39)
	Sharp Memorial Hospital	0.0% (0;3)	0.0% (0;5)	1.9% (1;31)	7.3% (8;113)	3.1% (11;289)	0.0% (0;52)	6.3% (24;426)	5.5% (31;643) Worse	13.0% (49;426) Worse	0.4% (1;309)	5.4% (11;252) Worse	4.4% (23;588)
	Tri-City Medical Center – Oceanside			5.5% (1;23)	7.3% (6;52)	1.7% (3;235)	0.0% (0;69)	5.9% (17;296)	3.6% (16;465)	13.6% (54;412) Worse	2.7% (9;306)	2.2% (4;211)	3.9% (19;488)
	UC San Diego Medical Center	0.0% (0;4)	5.8% (1;24)	0.0% (0;14)	4.0% (11;267)	1.2% (5;529)	3.9% (1;27)	2.7% (5;261) Better	2.6% (10;384)	9.0% (34;307)	1.2% (3;251)	3.4% (3;85)	2.8% (10;389)
San Francisco	California Pacific Medical Center – Pacific Campus	0.0% (0;3)	3.8% (2;38)	6.7% (1;17)	5.7% (16;200)	1.7% (10;305)	2.3% (1;40)	6.2% (16;231)	4.4% (29;529)	10.9% (80;515)	2.9% (11;310)	1.5% (4;150)	4.0% (15;322)
	California Pacific Medical Center – St. Luke’s Campus							9.6% (4;14)	7.0% (7;134)	24.2% (2;18)	9.4% (4;68) Worse	0.0% (0;29)	8.2% (12;132) Worse

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Chinese Hospital							3.4% (1;19)	3.7% (4;103)	5.2% (1;60)	4.0% (2;70)	0.0% (0;28)	7.1% (11;146)
	Kaiser Foundation Hospital – San Francisco		0.0% (0;19)	0.0% (0;47)		1.0% (7;919) Better	1.2% (1;50)	3.6% (15;775) Better	3.3% (12;380)	4.6% (10;272) Better	2.8% (4;202)	4.0% (3;76)	4.5% (10;185)
	Laguna Honda Hospital and Rehabilitation Center												3.4% (1;35)
	Saint Francis Memorial Hospital				8.3% (4;18)		0.0% (0;3)	11.2% (5;24)	4.4% (8;191)	9.8% (17;147)	2.0% (2;112)	2.3% (1;52)	8.3% (20;227) Worse
	Saint Mary’s Medical Center, San Francisco			0.0% (0;5)		2.6% (7;279)	0.0% (0;13)	9.6% (11;157)	2.9% (8;216)	6.5% (6;98)	1.3% (2;105)	2.3% (1;67)	4.1% (9;193)
	San Francisco General Hospital		0.0% (0;3)		16.0% (18;73) Worse	3.0% (3;93)		12.6% (13;159) Worse	5.5% (9;361)	16.7% (42;238) Worse	0.7% (1;184)	9.0% (2;33)	5.8% (9;327)
	UC San Francisco Medical Center	4.3% (1;28)	0.0% (0;34)	0.0% (0;39)	6.2% (51;1,266)	2.9% (8;150)	0.0% (0;28)	9.1% (14;143)	5.1% (18;370) Worse	11.6% (81;493)	2.5% (5;167)	4.4% (4;84)	5.3% (22;437)
San Joaquin	Dameron Hospital					5.2% (6;189) Worse	0.0% (0;7)	11.1% (12;235) Worse	6.4% (12;267) Worse	9.7% (7;120)	5.6% (8;155) Worse	9.5% (3;50)	4.8% (11;274)
	Doctors Hospital of Manteca							18.0% (4;14)	8.6% (7;108) Worse	5.0% (1;45)	2.5% (1;63)	0.0% (0;20)	3.7% (4;129)
	Kaiser Foundation Hospital – Manteca		0.0% (0;4)	0.0% (0;17)			0.0% (0;47)	8.2% (9;81)	4.2% (11;282)	4.6% (4;131)	3.4% (4;126)	0.0% (0;100)	3.7% (6;219)
	Lodi Memorial Hospital			0.0% (0;4)			0.0% (0;18)	8.4% (5;44)	1.3% (3;255)	12.8% (12;127)	3.3% (4;145)	5.5% (3;76)	3.1% (8;312)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Saint Joseph's Medical Center of Stockton	0.0% (0;3)		0.0% (0;26)	2.7% (2;60)	0.5% (2;516) Better	0.0% (0;89)	7.4% (23;371)	4.2% (23;523)	9.4% (26;280)	1.3% (4;295)	3.2% (5;139)	5.3% (28;511)
	San Joaquin General Hospital				11.3% (1;10)		0.0% (0;6)	6.9% (4;72)	5.0% (8;266)	10.2% (13;118)	3.6% (4;141)	0.0% (0;28)	4.2% (5;202)
	Sutter Tracy Community Hospital							16.8% (7;19) Worse	1.7% (2;108)	12.2% (4;49)	1.9% (1;64)	0.0% (0;30)	1.8% (2;140)
San Luis Obispo	Arroyo Grande Community Hospital						0.0% (0;14)	10.7% (5;22)	3.2% (2;99)	4.0% (1;58)	8.5% (5;92) Worse	6.6% (3;56)	1.8% (1;85)
	French Hospital Medical Center					1.6% (4;371)	0.0% (0;35)	10.2% (13;224)	2.2% (3;120)	9.0% (3;48)	3.2% (3;97)	1.9% (1;53)	3.4% (4;119)
	Sierra Vista Regional Medical Center			0.0% (0;17)	10.3% (7;89)		0.0% (0;18)	13.7% (4;14)	1.9% (1;69)	15.4% (25;143) Worse	0.0% (0;93)	2.9% (1;50)	8.3% (5;118)
	Twin Cities Community Hospital						0.0% (0;12)	11.1% (4;28)	6.3% (9;165)	32.0% (21;118) Worse	6.5% (6;157) Worse	4.8% (2;67)	6.7% (11;256)
San Mateo	Kaiser Foundation Hospital – Redwood City		0.0% (0;4)		7.6% (32;583)	0.0% (0;10)		6.3% (4;47)	5.8% (12;191) Worse	12.5% (62;357) Worse	2.3% (2;100)	1.8% (1;92)	7.2% (13;162)
	Kaiser Foundation Hospital – South San Francisco	0.0% (0;4)					0.0% (0;4)	4.3% (5;56)	4.9% (11;185)	12.5% (18;145)	3.2% (5;163)	1.9% (2;95)	3.4% (9;181)
	Peninsula Medical Center		0.0% (0;7)	0.0% (0;19)	10.2% (5;21)	1.7% (4;170)	0.0% (0;38)	6.4% (11;162)	2.6% (12;327)	13.7% (32;219) Worse	2.8% (8;227)	2.5% (5;194)	4.1% (15;332)
	San Mateo Medical Center				0.0% (0;7)		0.0% (0;4)	14.6% (7;29)	4.1% (2;91)	17.2% (7;45)	0.0% (0;68)	8.6% (2;18)	1.2% (1;91)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Sequoia Hospital			0.0% (0;18)	6.6% (1;7)	0.0% (0;90)	0.0% (0;57)	6.2% (5;87)	3.9% (10;255)	14.6% (15;103)	4.4% (4;86)	3.9% (2;71)	3.5% (5;153)
	Seton Medical Center			0.0% (0;7)	14.1% (5;22)	2.8% (11;269)	0.0% (0;24)	9.6% (15;164)	3.5% (15;437)	11.7% (30;222)	6.8% (10;160) Worse	0.0% (0;83)	5.2% (20;316)
Santa Barbara	Goleta Valley Cottage Hospital							27.6% (1;4)	0.0% (0;43)	13.8% (1;6)	7.5% (1;7)	7.1% (3;80)	5.6% (2;35)
	Lompoc Valley Medical Center						0.0% (0;7)	18.3% (2;11)	8.4% (5;70)	17.5% (3;27)	3.5% (1;59)	0.0% (0;31)	8.7% (10;121) Worse
	Marian Medical Center			0.0% (0;8)	0.0% (0;6)	2.7% (4;156)	0.0% (0;46)	9.9% (17;168)	3.2% (13;236)	14.6% (21;179)	2.0% (4;177)	2.2% (2;90)	4.8% (10;161)
	Santa Barbara Cottage Hospital	0.0% (0;3)	0.0% (0;7)	12.1% (2;26) Worse	5.8% (13;227)	1.8% (7;446)	0.0% (0;61)	10.1% (20;235)	3.5% (14;378)	10.7% (55;473)	1.2% (3;266)	4.9% (6;133)	4.5% (13;366)
	Santa Ynez Valley Cottage Hospital								10.2% (2;22)	0.0% (0;4)	0.0% (0;3)		6.7% (2;34)
Santa Clara	El Camino Hospital		0.0% (0;5)	3.9% (1;58)	4.5% (3;41)	2.7% (7;217)	0.0% (0;36)	7.5% (14;201)	2.9% (13;441)	12.1% (34;302)	2.2% (6;311)	3.5% (5;193)	6.9% (26;400) Worse
	Good Samaritan Hospital – San Jose		100% (2;3) Worse	0.0% (0;4)	9.9% (17;117)	2.8% (10;256)	0.0% (0;23)	4.7% (9;218)	5.2% (21;313) Worse	10.6% (38;318)	2.8% (7;207)	2.2% (4;148)	6.5% (19;258)
	Kaiser Foundation Hospital – San Jose		0.0% (0;4)	0.0% (0;6)		1.9% (1;58)	0.0% (0;26)	9.4% (17;157)	4.6% (16;346)	10.8% (27;294)	1.2% (2;267)	4.0% (5;113)	6.7% (20;334) Worse
	Kaiser Foundation Hospital – Santa Clara	0.0% (0;6)		0.0% (0;24)		2.5% (22;1,506)	0.0% (0;42)	8.2% (35;871)	3.5% (24;626)	10.9% (25;281)	1.9% (5;292)	4.7% (6;160)	4.8% (21;339)
	O'Connor Hospital – San Jose		0.0% (0;6)	0.0% (0;16)	0.0% (0;35)	1.4% (4;228)	0.0% (0;25)	4.1% (7;156)	3.4% (14;388)	11.3% (18;178)	2.2% (4;224)	3.6% (3;120)	5.0% (20;283)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Regional Medical of San Jose				8.1% (12;60)	1.6% (9;255)	2.2% (1;10)	6.7% (22;232)	3.1% (14;389)	11.1% (51;297)	1.8% (4;183)	3.3% (3;62)	6.3% (19;320)
	Saint Louise Regional Hospital							19.4% (4;18)	3.3% (5;135)	9.5% (3;32)	3.9% (2;67)	0.0% (0;35)	3.8% (4;127)
	Santa Clara Valley Medical Center				8.8% (5;78)	2.5% (3;172)	0.0% (0;9)	5.0% (8;215)	1.5% (4;523)	9.5% (20;297)	3.2% (6;259)	0.0% (0;46)	4.1% (13;409)
	Stanford Hospital	3.0% (1;27)	2.6% (1;37)	0.0% (0;73)	3.4% (25;764) Better	1.8% (6;247)	1.5% (1;79)	4.2% (10;208)	4.2% (28;465)	8.2% (52;450)	1.1% (5;329)	1.4% (2;131)	3.6% (18;428)
Santa Cruz	Dominican Hospital – Santa Cruz/Soquel				8.5% (4;59)	3.0% (11;289)	0.0% (0;31)	10.4% (17;208)	3.1% (8;309)	13.1% (19;212)	2.4% (4;254)	0.7% (1;115)	5.7% (12;271)
	Watsonville Community Hospital							6.1% (3;35)	5.3% (8;189)	8.3% (5;71)	4.0% (2;83)	0.0% (0;35)	4.6% (6;147)
Shasta	Mayers Memorial Hospital								24.1% (1;9)	67.3% (1;3)	0.0% (0;3)		20.7% (1;21)
	Mercy Medical Center – Redding			0.0% (0;33)	7.3% (5;79)	1.6% (9;375)	0.0% (0;124)	3.5% (11;334) Better	3.7% (18;344)	9.9% (33;281)	1.5% (5;188)	2.6% (7;187)	3.5% (16;383)
	Shasta Regional Medical Center					1.6% (4;239)	0.0% (0;44)	3.7% (7;171)	1.8% (9;301)	8.1% (10;177)	1.8% (6;187)	1.3% (2;68)	0.9% (4;304) Better
Siskiyou	Fairchild Medical Center						0.0% (0;5)	19.7% (1;4)	3.6% (1;28)	5.8% (1;23)	0.0% (0;44)	3.1% (1;25)	6.6% (6;128)
	Mercy Medical Center Mt. Shasta						0.0% (0;6)	0.0% (0;8)	0.0% (0;46)	26.6% (5;20) Worse	12.0% (2;11)	0.0% (0;12)	8.2% (4;60)
Solano	Kaiser Foundation Hospital – Rehabilitation Center Vallejo			4.6% (1;15)			0.0% (0;28)	4.5% (5;78)	3.8% (14;343)	13.7% (26;265)	3.3% (5;220)	6.3% (6;90) Worse	3.5% (14;377)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Kaiser Foundation Hospital – Vacaville							1.8% (1;46)	7.1% (12;162) Worse	19.4% (13;101) Worse	0.0% (0;75)	1.6% (1;58)	5.0% (12;264)
	North Bay Medical Center			0.0% (0;4)		1.7% (3;123)	3.2% (1;13)	8.1% (8;90)	3.9% (6;202)	11.6% (7;75)	2.2% (2;82)	3.2% (2;37)	5.1% (10;188)
	North Bay VacaValley Hospital							5.0% (2;20)	3.4% (4;124)	13.8% (7;53)	2.8% (2;68)	3.1% (1;32)	4.3% (8;141)
	Sutter Solano Medical Center						0.0% (0;5)	4.9% (2;24)	3.6% (6;177)	8.2% (5;104)	1.8% (2;90)	1.0% (1;35)	1.6% (2;126)
Sonoma													
	Healdsburg District Hospital							8.6% (1;6)	3.7% (1;33)	25.8% (6;29) Worse	5.2% (1;18)	0.0% (0;32)	2.8% (1;47)
	Kaiser Foundation Hospital – Santa Rosa						0.0% (0;34)	4.0% (4;90)	2.0% (3;161)	6.2% (8;198)	2.2% (3;157)	4.9% (3;92)	2.7% (5;212)
	Palm Drive Hospital						0.0% (0;5)		0.0% (0;28)	0.0% (0;28)	0.0% (0;29)	0.0% (0;31)	4.7% (3;70)
	Petaluma Valley Hospital						0.0% (0;4)	7.4% (7;42)	3.9% (4;83)	11.5% (5;38)	8.4% (4;61) Worse	1.4% (1;39)	9.4% (10;100) Worse
	Santa Rosa Memorial Hospital – Montgomery			0.0% (0;18)	12.8% (9;27) Worse	3.4% (7;208)	4.9% (1;35)	10.6% (22;231) Worse	5.7% (16;293) Worse	13.6% (37;256) Worse	2.6% (8;217)	4.4% (9;159)	5.2% (16;273)
	Sonoma Valley Hospital						0.0% (0;3)	11.3% (3;17)	4.2% (4;64)	9.9% (5;46)	2.6% (1;35)	3.4% (1;37)	6.4% (3;57)
	Sutter Medical Center of Santa Rosa			0.0% (0;11)		2.0% (3;223)	1.8% (1;43)	3.6% (4;184)	5.2% (5;132)	12.1% (3;42)	0.0% (0;80)	4.2% (2;34)	3.1% (2;97)
Stanislaus													
	Doctors Medical Center		0.0% (0;3)	3.5% (1;25)	7.5% (4;40)	1.6% (12;500)	1.5% (2;119)	7.4% (29;467)	7.3% (26;413) Worse	12.9% (41;307)	4.0% (8;194)	2.3% (2;115)	4.8% (13;326)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Emanuel Medical Center, Inc.			0.0% (0;6)			2.5% (1;36)	6.8% (9;65)	4.8% (16;298)	9.5% (8;126)	4.7% (7;221)	2.4% (2;76)	8.6% (28;383) Worse
	Memorial Hospital Medical Center – Modesto		22.0% (2;6)	0.0% (0;22)	6.8% (3;26)	2.7% (20;593)	0.5% (1;153)	7.2% (32;436)	3.7% (27;641)	9.0% (34;357)	1.6% (6;328)	2.0% (6;223)	4.8% (31;594)
	Oak Valley District Hospital							8.6% (2;12)	2.2% (1;47)	31.9% (6;31) Worse	8.7% (2;27)	0.0% (0;16)	3.4% (2;60)
Tehama	Saint Elizabeth Community Hospital							6.9% (2;21)	0.0% (0;93)	8.2% (4;63)	0.0% (0;64)	0.0% (0;33)	1.2% (1;128)
Trinity	Trinity Hospital							6.1% (1;9)	0.0% (0;14)	0.0% (0;5)	0.0% (0;7)		3.5% (1;51)
Tulare	Kaweah Delta Medical Center			0.0% (0;19)		2.1% (17;1,289)	2.1% (1;87)	9.4% (36;489) Worse	5.0% (19;458) Worse	15.4% (33;330) Worse	4.9% (12;311) Worse	6.5% (10;170) Worse	3.6% (19;662)
	Sierra View District Hospital							10.7% (6;52)	4.4% (12;260)	18.3% (17;111) Worse	3.7% (4;116)	0.0% (0;50)	8.7% (31;373) Worse
	Tulare Regional Medical Center							9.0% (4;28)	2.5% (2;128)	17.5% (6;56)	6.0% (2;58)	4.0% (2;39)	9.9% (15;239) Worse
Tuolumne	Sonora Regional Medical Center – Greenley						0.0% (0;36)	3.2% (2;21)	2.5% (5;91)	8.5% (6;64)	1.5% (2;98)	4.1% (4;72)	1.9% (7;226) Better
Ventura	Community Memorial Hospital – San Buenaventura			0.0% (0;14)	14.6% (3;27)	1.5% (4;361)	0.0% (0;48)	4.7% (11;258)	3.0% (11;351)	10.1% (18;228)	0.9% (2;212)	2.3% (3;124)	4.6% (13;262)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.

California Hospital Risk-Adjusted Mortality Rates (RAMR), Total Deaths and Cases, and Quality Ratings for Inpatient Mortality Indicators, 2010

County	Hospital Name	Esophageal Resection	Pancreatic Resection	Abdominal Aortic Aneurism Repair	Craniotomy	Percutaneous Coronary Intervention	Carotid Endarterectomy	Acute Myocardial Infarction	Heart Failure	Acute Stroke	Gastro-Intestinal Hemorrhage	Hip Fracture	Pneumonia
NOTE: In each cell, the first line indicates Risk-Adjusted Mortality Rate (RAMR), the second line indicates total deaths; total cases, and the third line indicates any quality rating.													
	STATEWIDE	3.9% (16;410)	4.5% (33;728)	1.4% (36;2,518)	6.6% (953;14,380)	2% (965;48,017)	0.5% (33;6,962)	6.8% (3,099;45,895)	3% (2,640;87,061)	10% (5,250;52,480)	2.2% (1063;48,089)	2.5% (584;23,740)	4.1% (3,108;75,063)
	Los Robles Hospital and Medical Center			0.0% (0;16)	9.9% (5;34)	1.6% (7;518)	0.0% (0;24)	8.1% (19;331)	4.2% (14;320)	11.9% (20;176)	3.5% (7;194)	3.1% (3;144)	5.3% (13;265)
	Ojai Valley Community Hospital							16.5% (2;4)	8.1% (1;19)	15.6% (5;28)	7.5% (1;19)	0.0% (0;27)	0.0% (0;56)
	Saint John's Pleasant Valley Hospital							7.8% (3;26)	3.1% (6;150)	9.0% (4;78)	0.8% (1;99)	3.8% (3;67)	3.5% (6;135)
	Saint John's Regional Medical Center			0.0% (0;18)	0.0% (0;24)	2.1% (13;287)	0.0% (0;35)	5.1% (22;298)	2.9% (14;409)	8.5% (24;202)	1.0% (3;204)	0.7% (1;93)	4.4% (17;303)
	Simi Valley Hospital and Healthcare Services – Sycamore				0.0% (0;12)		0.0% (0;6)	4.9% (5;51)	2.1% (3;133)	7.2% (7;114)	0.0% (0;90)	1.4% (1;63)	4.4% (7;169)
	Thousand Oaks Surgical Hospital				22.2% (2;51)								
	Ventura County Medical Center				6.1% (3;33)			11.7% (3;25)	2.3% (2;115)	7.6% (7;114)	1.2% (1;114)	4.2% (1;37)	5.6% (6;146)
Yolo	Sutter Davis Hospital							11.0% (2;5)	2.5% (2;64)	0.0% (0;23)	1.2% (1;60)	3.4% (2;52)	5.5% (5;97)
	Woodland Memorial Hospital						0.0% (0;18)	2.6% (1;27)	0.9% (1;56)	23.3% (11;74) Worse	0.0% (0;68)	1.3% (1;52)	1.9% (3;108)
Yuba	Rideout Memorial Hospital			0.0% (0;11)	4.1% (1;36)	2.6% (5;162)	0.0% (0;63)	6.8% (14;240)	4.2% (16;422)	12.0% (26;234)	1.4% (3;200)	5.5% (4;97)	3.9% (13;513)

Source: Agency for Healthcare Research and Quality (AHRQ) Quality Indicators and SAS, Version 4.4, and OSHPD hospital patient discharge data, 2010. See Technical Note for details about methodology.

*Hospital consolidated or changed name from 2010 to 2011, see OSHPD Implementation of AHRQ Inpatient Mortality Indicators, 2010 and 2011 data, Technical Note.

Note: Blank cells indicate that fewer than three procedures were performed or conditions were treated.

Definition of Terms: **Risk Adjustment:** The Risk Adjusted Mortality Rates (RAMR) presented here adjust the observed mortality rates. This statistical methodology takes into account pre-existing health problems that put some patients at greater risk of death to 'level the playing field' and allow fair comparisons across hospitals. **Total Deaths:** number of patients that died in this hospital. **Total Cases:** number of patients undergoing this procedure or with this medical condition in this hospital.

Quality Ratings are risk-adjusted using age, gender, APD-DRG base category, and APR-DRG risk of mortality scores. Significantly better or significantly worse than the statewide rate are noted when upper 95% confidence level of the RAMR falls below the statewide rate ('Better') or lower 95% confidence level of the RAMR is higher than the statewide rate ('Worse'). Cells with no quality ratings indicate a hospital's performance is considered 'not significantly different' from the statewide average.